

XP
H947
V.76
H6

PHYTOLOGIA

An international journal to expedite plant systematic, phytogeographical
and ecological publication

Vol. 76

June 1994

No. 6

CONTENTS

PALACIOS C., R. & D.L. QUIROZ G., Estudio palinologico de las especies Mexicanas del género <i>Stenandrium</i> Nees (Acanthaceae).	433
KARTESZ, J.T. & K.N. GANDHI, Nomenclatural notes for the North American flora XIII.	441
MARSH, J.E. & T.H. NASH, III, A new lichen species, <i>Niebla cedrosensis</i> (Ramalinaceae), is described from Baja California, México.	458
BABU, G.J., V. PUSHPALATHA, & K.C. NAIDU, Weed flora of Khammam District in Andhra Pradesh, India.	461
NESOM, G.L., A new species of <i>Castilleja</i> (Scrophulariaceae) related to <i>C. aspera</i> from southeastern Durango, México.	465
NESOM, G.L., Two new species of <i>Sisyrinchium</i> subg. <i>Echthronema</i> (Iridaceae) from Nuevo León, México.	468
SIVINSKI, R., T. LOWREY, & R. PETERSON, Additions to the native and adventive flora of New Mexico.	473
Books received.	480
Index to authors, volume 76.	482
Index to taxa, volume 76.	483
Index to reviewers, volume 76.	509
Corrections and additions.	511
Back issues available.	512

LIBRARY

DEC. 7 1994

NEW YORK
BOTANICAL GARDEN

Published by Michael J. Warnock
185 Westridge Drive Huntsville, Texas 77340 U.S.A.
PHYTOLOGIA is printed on acid free paper.

PHYTOLOGIA (ISSN 00319430) is published monthly with two volumes per year by Michael J. Warnock, 185 Westridge Drive, Huntsville, TX 77340. Second Class postage paid at Huntsville, TX. Copyright 1994 by PHYTOLOGIA. Annual domestic individual subscription (12 issues): \$40.00. Annual domestic institutional subscription (12 issues): \$44.00. Foreign and/or airmail postage extra. Single copy sales: Current issue and back issues volume 72 to present, \$4.00; Back issues (previous to volume 72), \$3.00 (add \$.75 per copy postage and handling US [\$1.50 per copy foreign]). Back issue sales by volume: \$17.00 per volume 42-71 (not all available as complete volumes); \$21.00 per volume 72-present; add \$3.00 per volume postage US (\$6.00 per volume foreign). POSTMASTER: Send address changes to Phytologia, 185 Westridge Drive, Huntsville, TX 77340-8916.

ESTUDIO PALINOLOGICO DE LAS ESPECIES MEXICANAS DEL GENERO *STENANDRIUM* NEES (ACANTHACEAE)

Rodolfo Palacios Chávez¹ & David Leonor Quiroz García¹

Escuela Nacional de Ciencias Biológicas del I.P.N., Prolongación Carpio y
Plan de Ayala, 11340 México, D.F., MEXICO

ABSTRACT

The pollen of seven Mexican species of *Stenandrium* is studied with light microscopy. The species are: *S. barbatum* Torrey & Gray, *S. chameranthemoideum* Oersted, *S. dulce* (Cav.) Nees, *S. manchonense* T.F. Daniel, *S. nanum* (Standley) T.F. Daniel, *S. pedunculatum* (Donn.-Smith) Leonard, and *S. pilosulum* (S.F. Blake) T.F. Daniel.

Pollen is usually prolate and tricolpate with differences in the ends of the colpi, and the ornamentation observed is faintly reticulate, punctitegillate, psilate or verrucate and gemmate.

Only *Stenandrium dulce* has pollen very different, not similar to the other species, being spherical, trichotomocolpate, verrucate and gemmate.

KEY WORDS: Palynology, *Stenandrium*, México, Acanthaceae

RESUMEN

El polen de siete especies mexicanas de *Stenandrium* es estudiado con el microscopio de luz; las especies corresponden a *S. barbatum* Torrey & Gray., *S. chameranthemoideum* Oersted, *S. dulce* (Cav.) Nees, *S. manchonense* T.F. Daniel, *S. nanum* (Standley) T.F. Daniel, *S. pedunculatum* (Donn.-Smith) Leonard, y *S. pilosulum* (S.F. Blake) T.F. Daniel.

El polen en general resulto ser prolato y tricolpado con diferencias en las terminaciones de los colpos y la ornamentación observada es ligeramente reticulada, puntitegilada, psilada o verrugada y gemada.

Únicamente *Stenandrium dulce* tiene polen muy diferente sin ninguna relación con el de las demás especies, el cual es esférico, tricotomocolpado, verrugado y gemado.

¹Becarios de la Comisión de Operación y Fomento de Actividades Académicas.

INTRODUCCION

Stenandrium es un género constituido por plantas herbáceas y subarborescentes que progresa en las regiones tropicales y subtropicales.

Aproximadamente 50 especies de este taxon han sido descritas para el continente americano (Daniel 1984) y tiene una distribución que abarca desde el sur de los Estados Unidos hasta Argentina y Chile. En territorio mexicano indica (Daniel 1984) que existen nueve especies: *Stenandrium barbatum* Torrey & Gray.; *S. chamerantheroideum* Oersted, *S. dulce* (Cav.) Nees, *S. manchonense* T.F. Daniel; *S. nanum* (Standley) T.F. Daniel; *S. pedunculatum* (Donn.-Smith) Leonard; *S. pilosulum* (S.F. Blake) T.F. Daniel; *S. subcordatum* Standley, y *S. verticillatum* Brandegees.

Desde el punto de vista palinológico, este género ha sido estudiado por Raj (1961) quien describe brevemente el polen de *Stenandrium barbatum* de Texas, como tricolpado, prolato ($48 \times 30 \mu$), colpos de $45 \times 4 \mu$ con terminaciones redondeadas; exina tectada de 3.5μ de grosor, densamente soportada por delgados báculos. El de *S. droseroides* de Cuba, tricolpado, prolato de ($35 \times 20 \mu$), finamente reticulado con apocolpio muy pequeño. El de *S. pohlii* de Brasil, pantoporado, esferoidal con diámetro de 40μ y con la exina verrugada. El de *S. trinerve* (Uruguay) con aberturas irregulares, esferoidal cerca de 32μ de diámetro, con la exina verrugada y finalmente el de *S. dulce* descrito como tectado, con el tegilum ondulado, soportado por delgados báculos.

Heusser (1971) describe el polen de *Stenandrium dulce* de la flora de Chile como inaperturado, verrugado a gemado, esferoidal de 30μ de diámetro. Markgraf & D'Antoni (1978) hacen la descripción de *S. cf. trinerve* como inaperturado, verrugado a gemado, esferoidal de 30μ de diámetro.

El trabajo más completo que se ha desarrollado sobre este taxon en el área de la palinología es el de Furness (1993) quien estudia el polen de catorce especies procedentes de Africa y Madagascar con microscopio de luz, barrido y transmisión. Encontrando que el polen de este género tiene suficientes variaciones para establecer cuatro tipos de polen a mencionar:

I. Polen generalmente prolato, tricolpado, con colpos sin márgenes prominentes, reticulado, perforado o fosulado.

II. Oblato esferoidal, 3-6 colpado con aberturas irregulares, finamente reticulado.

III. Perprolato, tricolpado, colpos con márgenes prominentes, finamente rugulado o perforado.

IV. Prolato a perprolato, tricolpado con márgenes prominentes y puentes mesocolpiales, y la ornamentación variable.

MATERIALES Y METODOS

El material de estudio (muestras florales) fue obtenido del herbario de la Escuela Nacional de Ciencias Biológicas del Instituto Politécnico Nacional (ENCB) a excepción de las especies de *Stenandrium nanum* y *S. pedunculatum* que se obtuvieron del herbario del Instituto de Biología de la Universidad Nacional Autónoma de México (MEXU). No se incluyen las descripciones de los granos de polen de *S. subcordatum* y de *S. verticillatum* porque no fue posible obtener muestras de estas especies.

El material floral obtenido fue tratado por medio de la técnica de acetolisis de Erdtman (1943) y las observaciones de los granos de polen se realizaron mediante el microscopio de luz.

DESCRIPCION DE LOS GRANOS DE POLEN

Stenandrium barbatum Torrey & Gray. 4 km al SW de Ciudad Juárez, Chihuahua. G. Borja 400 (ENCB). Lám. I. Figs. 1 a 4.

Polen tricolpado, tectado, prolato, de $51(56)60 \times 35(37)40 \mu$. P/E=1.5. Vista polar circular de $38(40)44 \mu$ de diámetro. Exina de 2.4μ de grosor, sexina de 1.6μ y la nexina de 0.8μ de espesor, superficialmente puntitegilada. Colpos cubiertos con membranas lisas y con terminaciones circulares.

Stenandrium chamerantheoideum Oerst. Pachuquilla, Mun. de Puente Nacional, Veracruz. F. Ventura 7764 (ENCB). Lám. I. Figs. 5 a 8.

Polen tricolpado, semitectado, prolato, de $42(47)53 \times 28(31)34 \mu$. P/E=1.5. Vista polar circular de $32(36)40 \mu$ de diámetro. Exina de 2.4μ de grosor, sexina de 1.6μ y nexina de 0.8μ de espesor, superficie finamente reticulada. Colpos operculados y con terminaciones circulares. Opérculo central finamente reticulado, la situación de esta estructura a veces es confusa, de tal forma que algunos granos dan la impresión de ser hexacolpados.


Stenandrium dulce (Cav.) Nees. Cerro Gordo, Edo. de México. J. Rzedowski 22205 (ENCB). Lám. I. Figs. 9 a 11.

Polen tricotomocolpado, intectado, esférico, de $43.0(45.0)51.2 \mu$ de diámetro. Exina de 5.6μ de grosor, con la sexina de 3.2μ y la nexina de 2.4μ de espesor, superficialmente verrugada y gemada, gemas de 4 a 5μ de diámetro y las verrugas de $3(5)6 \mu$ de alto \times $7(8)12 \mu$ de ancho. Colpo en forma de "Y" con las membranas escabrosas.

Stenandrium manchonense T.F. Daniel. Manchón, Distrito de Mina, Guerrero. Hinton 10460 (ENCB). Lám. I. Figs. 12 a 15.

Polen dicolpado, tectado, prolato, de $41(47)52 \times 22(27)34 \mu$. P/E=1.77. Vista polar cuadrangular de $28(33)38 \mu$ de diámetro.

Exina de 1.8μ de grosor, con la sexina de mayor espesor que la nexina, superficialmente puntitegilada. Colpos con los márgenes irregulares cubiertos con membranas lisas y con terminaciones circulares.



LAMINA I

Lámina I. *Stenandrium barbatum*: 1. Sección óptica de la vista ecuatorial; 2. Vista ecuatorial superficial; 3. Vista polar en sección óptica; 4. Vista polar mostrando las terminaciones de los colpos. *S. chameranthemoideum*: 5. Vista ecuatorial en sección óptica; 6. Vista ecuatorial superficial; 7. Vista polar mostrando el grosor de la exina; 8. Vista polar mostrando la ornamentación. *S. dulce*: 9. Sección óptica. 10. Abertura trirradiada y ornamentación; 11. Vista superficial. *S. manchonense*: 12. Vista ecuatorial en sección óptica; 13. Vista ecuatorial superficial; 14. Vista polar en sección óptica; 15. Vista polar mostrando las terminaciones de los colpos.

Stenandrium nanum (Standley) T.F. Daniel. Rancho de dos Fierros, 7 km al S de la desviación a Puerto Morelos, Quintana Roo. *E.F. Cabrera 6313* (MEXU). Lám. II. Figs. 16 a 19.

Polen tricolpado, semitectado, prolato, de $46(54)61 \times 24(31)37 \mu$. P/E=1.74. No se pudo observar vistas polares. Exina de 1.6μ de grosor, con la sexina y la nexina de igual espesor, superficie ligeramente reticulada. Colpos cubiertos con membranas escabrosas y con las terminaciones agudas.

Stenandrium pedunculatum (Donn.-Smith) Leonard. Estación de Biología de Chamela, Jalisco. *E. Lott 1366* (MEXU). Lám. II. Figs. 20 a 22.

Polen tricolpado, tectado, prolato, de $41(52)63 \times 23(30)44 \mu$. P/E=1.7. Vista polar circular de $24(32)46 \mu$ de diámetro. Exina de 3.2μ de grosor, con la sexina y la nexina de igual espesor, superficie ligeramente reticulada. Colpos cubiertos con membranas granulosas, abriéndose hacia los extremos con terminaciones circulares, que aparentan ser poros cubiertos con membranas escabrosas.

Stenandrium pilosulum (S.F. Blake) T.F. Daniel. 7 millas al NW de Yecare, Sonora. *R. Moran 21965* (ENCB). Lám. II. Figs. 23 a 27.


Polen dicolpado, algunas veces tricolpado, tectado, prolato, de $50(54)62 \times 32(38)43 \mu$. P/E=1.5. Vista polar circular de $38(40)44 \mu$ de diámetro. Exina de 2.4μ de grosor, sexina de 1.6μ y la nexina de 0.8μ de espesor, superficialmente psilada. Colpos constreñidos en el ecuador, abriéndose hacia los extremos en forma de poros circulares, cubiertos con membranas granulosas.

CONCLUSIONES

El polen de las especies mexicanas del género *Stenandrium* resulto ser prolato en la mayor parte de los taxa en estudio, a excepción de *S. dulce* que lo tiene esférico, sin embargo otras diferencias observadas radican principalmente en las aberturas y en la ornamentación de esos granos de polen. Con referencia a la primera observación es notable que algunas especies tienen colpos con terminaciones circulares dando la impresión que poseen poros en los extremos de los colpos, es dicolpado con estas características en *S. barbatum* y tricolpado en *S. manchonense*, *S. pedunculatum*, y *S. pilosulum*. Polen con otro tipo de aberturas se aprecia en *S. chameranthemoideum* donde es tricolpado y los colpos son operculados de tal forma que algunos granos dan la impresión de ser hexacolpados. Tricolpado con terminaciones agudas fue encontrado en *S. nanum*. Tricotomocolpado sólo fue observado en *S. dulce*.

En lo que a ornamentación se refiere se puede apreciar que esta es poco conspicua, resulto ser puntitegilada en el polen de *Stenandrium barbatum* y *S. manchonense*; psilada en *S. pilosulum* y ligeramente reticulada en *S. chameranthemoideum*, *S. nanum*, y *S. pedunculatum*.

Comparando los resultados obtenidos con los de otros autores el polen de *Stenandrium barbatum* coincide con las características dadas por Raj (1961),



LAMINA II

Lámina II. *Stenandrium nanum*: 16. Vista ecuatorial en corte óptico; 17. Vista ecuatorial superficial mostrando las aberturas; 18. Vista ecuatorial superficial; 19. Vista ecuatorial donde se aprecia uno de los colpos. *S. pedunculatum*: 20. Vista ecuatorial sección óptica. 21. Vista ecuatorial mostrando uno de los colpos; 22. Vista ecuatorial superficial. *S. pilosulum*: 23. Vista ecuatorial en sección óptica; 24. Vista ecuatorial mostrando dos de los colpos; 25. Vista ecuatorial superficial; 26. Contorno de la vista polar; 27. Vista polar donde se aprecia la terminación de los colpos.

Tabla 1. Principales características palinológicas establecidas en las especies del género *Stenandrium*.

Especie	Forma	Abertura	Terminación de los colpos	Ornamentación
<i>Stenandrium barbatum</i>	Prolato	Tricolpado	Circular	Puntitegilada
<i>S. chameranthemoideum</i>	Prolato	Tricolpado	Circular	Reticulada
<i>S. dulce</i>	Esférico	Tricotomo colpado	—	Verrugada y gemada
<i>S. manchonense</i>	Prolato	Dicolpado	Circular	Puntitegilada
<i>S. nanum</i>	Prolato	Tricolpado	Aguda	Reticulada
<i>S. pedunculatum</i>	Prolato	Tricolpado	Circular	Reticulada
<i>S. pilosulum</i>	Prolato	Tricolpado	Circular	Psilada

pero para el de *S. dulce* son diferentes en lo referente a tectum y ornamentación, por lo que quizás se trate de otro taxon. De acuerdo con los tipos establecidos por Furness (1993) el polen de las especies *S. barbatum* y *S. nanum* se pueden situar dentro del tipo I; polen oblato no se pudo observar, tampoco el esferoidal, el perprolato, el margocolpado, ni con puentes mesocolpiales perteneciente a los otros grupos que señala el mismo autor.

Es indudable que las especies mexicanas de *Stenandrium* tienen polen muy diferente al de las especies de África y Madagascar. Sobresale el polen de *S. dulce* que por sus características se aleja por completo de cualesquiera de las especies que aquí se mencionan.

Las diferencias palinológicas más notorias que podemos señalar en el género *Stenandrium* se resumen en la Tabla 1.

LITERATURA CITADA

- Daniel, T.F. 1984. A revision of *Stenandrium* (Acanthaceae) in Mexico and adjacent regions. Ann. Missouri Bot. Gard. 71:1028-1043.
- Erdtman, G. 1943. *An Introduction to Pollen Analysis*. The Ronald Press Company, New York, New York. 239 pp.
- Furness, C.A. 1993. A pollen morphological survey of the old world species of *Stenandrium* Nees (Acanthaceae: Acantheae). Grana 32:1-11.
- Heusser, C.J. 1971. *Pollen and Spores of Chile*. University of Arizona Press, Tucson, Arizona 167 pp.

Markgraf, V. & H.L. D'Antoni. 1978. *Pollen Flora of Argentina*. The University of Arizona Press, Tucson, Arizona. 143 pp.

Raj, B. 1961. Pollen morphological studies in the Acanthaceae. *Grana Palynologica* 3:3-108.

NOMENCLATURAL NOTES FOR THE NORTH AMERICAN FLORA. XIII.

John T. Kartesz & Kancheepuram N. Gandhi

North Carolina Botanical Garden, Dept. of Biology, Coker Hall, University of North Carolina, Chapel Hill, North Carolina 27599-3280, U.S.A.

ABSTRACT

The nomenclature and taxonomy of the following are discussed: *Blastania*, *Caltha palustris* var. *flabellifolia*, *Chaenomeles*, *Choenomeles*, *Ctenolepis*, *Heracleum lanatum*, *H. maximum*, *Coincya monensis* var. *recurvata*, *Frangula*, *Laburnum*, *Physalis izocarpa*, *Physalis philadelphica*, *Prunus pumila* vars. *cuneata* and *susquehanae*, *Ranunculus glaberimus* var. *reconditus*, *Rhamnus*, *Rhamnus davurica* and *Rhamnus citrifolia*. Fourteen new combinations and one *status novum* are proposed: **Frangula betulifolia** (E. Greene) V. Grub. ssp. *obovata* (Kearney & Peebles) Kartesz & Gandhi, *comb. nov.*; **Frangula** × **blumeri** (E. Greene) Kartesz & Gandhi, *comb. nov.*; **Frangula californica** (Eschsch.) A. Gray ssp. *crassifolia* (Jepson) Kartesz & Gandhi, *comb. nov.*; **Frangula californica** ssp. *cuspidata* (E. Greene) Kartesz & Gandhi, *comb. nov.*; **Frangula californica** ssp. *occidentalis* (T.J. Howell) Kartesz & Gandhi, *comb. nov.*; **Frangula californica** ssp. *toментella* (Benth.) Kartesz & Gandhi, *comb. nov.*; **Frangula californica** ssp. *ursina* (E. Greene) Kartesz & Gandhi, *comb. nov.*; **Frangula rubra** (E. Greene) V. Grub. ssp. *modocensis* (C.B. Wolf) Kartesz & Gandhi, *comb. nov.*; **Frangula rubra** ssp. *nevadensis* (A. Nels.) Kartesz & Gandhi, *comb. nov.*; **Frangula rubra** ssp. *obtusissima* (E. Greene) Kartesz & Gandhi, *comb. nov.*; **Frangula rubra** ssp. *yosemitana* (C.B. Wolf) Kartesz & Gandhi, *comb. nov.*; **Frangula sphaerosperma** (Sw.) Kartesz & Gandhi, *comb. nov.*; **Frangula sphaerosperma** ssp. *longipes* (M.C. & L.A. Johnston) Kartesz & Gandhi, *comb. nov.*; **Physalis ixocarpa** Brot. ex Hornem. var. *immaculata* (Waterfall) Kartesz & Gandhi, *comb. nov.* and var. *parviflora* (Waterfall) Kartesz & Gandhi, *comb. nov.*; **Rhamnus davurica** Pallas ssp. *nipponica* (Makino) Kartesz & Gandhi, *comb. nov.*; **R. lanceolata** Pursh ssp. *glabrata* (Gleason) Kartesz & Gandhi, *stat. nov.*

KEY WORDS: varietal autonyms, Apiaceae, Brassicaceae, Cucurbitaceae, Fabaceae, Ranunculaceae, Rhamnaceae, Rosaceae, Solanaceae, *Blastania*, *Coincya*, *Caltha*, *Chaenomeles*, *Choenomeles*, *Ctenolepis*, *Frangula*, *Heracleum*, *Laburnum*, *Physalis*, *Prunus*, *Ranunculus*, *Rhamnus*

INTRODUCTION

Continuing with the "NOMENCLATORIAL NOTES FOR THE NORTH AMERICAN FLORA" (Kartesz & Gandhi 1989; 1990a, b, c; 1991a, b, c, d; 1992a, b, c, d), a thirteenth note in the series is presented here toward advancing our understanding of North American plants.

APIACEAE

Heracleum mazimum and *H. lanatum*

Contrary to Fernald's (1950, p. 1104) and Voss' (1985, p. 661) treatments of the Cow-parsnip as *Heracleum mazimum* Bartr. (published in 1791), Gleason & Cronquist (1991, p. 382) accepted the name *H. lanatum* Michx. (published in 1803) for this species and cited *H. mazimum* as its synonym. Regarding the usage of one name over another, the dispute is not over the taxonomy, nor over name priority, but rather over the validity of Bartram's binomials published in his 1791 work. A brief discussion is here provided.

Bartram (1791), in his work titled "*Travels through North and South Carolina . . .*", published 49 new binomials (Wilbur 1971). At the same time he also used a few polynomials (e.g., p. 378, "*Pinus palustris, foliis trinis longissimis, strobilo elongato . . .*"). Due to his occasional use of polynomials, the validity of the 49 binomials has been in dispute. Based on Art. 68.4 of the 1935 Code (see Camp 1947; presently *International Code of Botanical Nomenclature (ICBN)* Art. 23.6c; Greuter 1988), Rickett (1944) rejected Bartram's binomials. However, Merrill (1945) and Wilbur (1971) stated that such occasional usage of polynomials are also found in Linnaeus (1753; e.g., p. 213: *Apocynum fol. androesimifolium*) and in P. Miller (1768; e.g., *Alchemilla foliis lobatis sericeis acutis, Aloe foliis erectis subulatis radicatis undique inerme spinosis, Asparagus caule herbaceo erecto, foliis setaceis*).

Both Merrill (1945) and Wilbur (1971) argued that Bartram's usage of polynomials was an exception rather than a rule, and therefore, his new binomials should be treated as valid. Wilbur listed Bartram's 49 new binomials. We concur with Merrill and Wilbur, and conclude that the name *Heracleum mazimum* was validly published in Bartram's work.

Heracleum maximum Bartr., *Travels*. 344. 1791.

SY: *Heracleum lanatum* Michx., *Fl. Bor. Amer.* 1:166. 1803.

BRASSICACEAE

Coincya monensis var. *recurvata*

Coincya monensis (L.) Greuter & Burdet, a European native, has become established in Pennsylvania, and New Jersey, and southward to North Carolina. The North American expression of this species is referable to var. *recurvata* (Rollins 1993, p. 329). Regarding the nomenclature of this variety, our analysis follows.

Leadlay & Heywood (1990, pp. 370-380) recognized four varieties within the protologue of the new combination *Coincya monensis* ssp. *recurvata* (All.) Leadlay. They (p. 375) did not cite authorship for var. *recurvata*. It appears as though these authors treated var. *recurvata* as a typical expression of ssp. *recurvata* and considered that authorship was unnecessary for the varietal autonym. Although permitted prior to the Sydney Congress, this practice is unacceptable under the present *Code*, i.e., the name *Coincya monensis* var. *recurvata* represents an autonym of a non-typical subspecies.

We (Kartesz & Gandhi, *Taxon* 40:308-310. 1991.) argued that varietal autonoms of non-typical subspecies must be treated as validly published, which was opposed (see *Taxon* 41:158. 1992.). We (*Taxon* 41:784. 1993.) countered the opposition by proposing that a new Art. 32.9, along with an example be added to the *Code*. The Tokyo Congress referred the example to the Editorial Committee of the *ICBN* for consideration (see *Taxon* 42:916. 1994.).

In light of the above situation, we consider the varietal autonym *Coincya monensis* var. *recurvata* to be validly published.

Coincya monensis (L.) Greuter & Burdet ssp. *recurvata* (All.) Leadlay, J. Linn. Soc. Bot. 102:370. 1990. BASIONYM: *Synapsis recurvata* All., *Fl. Pedem.* 1:2654. 1785.

Coincya monensis (L.) Greuter & Burdet var. *recurvata* (All.) Leadlay, J. Linn. Soc. Bot. 102:375. 1990. BASIONYM: *Synapsis recurvata* All., *Fl. Pedem.* 1:2654. 1785.

CUCURBITACEAE

Blastania, *Ctenolepis*

Ctenolepis cerasiformis (Stocks) Hook.f., a native of tropical Africa, has been reported for Maryland. For this taxon, Meeuse (1962, p. 12) used the name *Blastania cerasiformis* (Stocks) A. Meeuse. In our study, we found that the genus *Blastania* Kotschy & Peyritsch has been treated as a taxonomic synonym of *Ctenolepis* Hook.f. (Hutchinson 1954; Welman 1987, p. 201) or vice versa (Meeuse, p. 11). Our analysis follows.

Regarding the effective publication of the genus *Blastania* Kotschy & Peyritsch, Jackson (1895) and Meeuse (1962, p. 11) mentioned the date to be 1865-66, whereas C. Jeffrey (KEW; pers. comm.) believes the date to be Oct 1867. However, we concur with Stafleu & Cowan (1979), who stated the publication date of Kotschy & Peyritsch's work to be Jul 1867.

For *Ctenopsis* Naudin (published in 1866, *non de Notaris* 1847), Hooker (in Bentham & Hooker, Sep 1867, p. 832) published *Ctenolepis* Hook.f. as a substitute name. With this analysis, we assert that the name *Blastania* is older than the name *Ctenolepis*.

Since *Blastania* and *Ctenolepis* represent the same entity and since *Blastania* has priority over *Ctenolepis*, the former should be accepted to be the correct name. To date, *Ctenolepis* has not been conserved over *Blastania*. In this situation, *Ctenolepis* should be treated as a heterotypic taxonomic synonym of *Blastania*.

Blastania cerasiformis (Stocks) A. Meeuse, *Bothalia* 8:12. 1962. BASIONYM: *Zehneria cerasiformis* Stocks, *J. Bot. Kew Gard. Misc.* 4:149. 1852. SY: *Ctenolepis cerasiformis* (Stocks) Hook.f., *Fl. Trop. Africa* 1:178. 1871.

FABACEAE

Laburnum

The Golden-chain tree (*Laburnum anagyroides* Medik.), a European native and extensively used in horticultural plantings in the eastern United States, has been reported as an escape in Massachusetts and elsewhere. The name *Laburnum anagyroides* was based on *Cytisus laburnum* L. The genus name *Laburnum* has been attributed either to Fabricius (Frodin & Heywood in Tutin, et al. 1968; Greuter 1989; Wiersema 1990; Brummitt 1993) or to Medikus (Jackson 1895).

Within Fabaceae, Fabricius (1759) proposed the genus name *Laburnum*, but did not designate a type. Twenty eight years later, Medikus (see below) proposed *Laburnum* as a new genus within Fabaceae. Pfeiffer (*Nom.* 2[1]:1874) lectotypified *Laburnum* Medik. by *L. anagyroides* Medik. All those

who recognize *Laburnum* Fabr. include *L. anagyroides* within it. Regarding the nomenclature of *Laburnum*, our analysis is given below.

If the name *Laburnum* Fabr. was validly published, then *Laburnum* Medik. is a later homonym. *Laburnum* Medik. is not an isonym, since its type is not that of *Laburnum* Fabr. (To the best of our knowledge, the latter has not been typified.). Dandy (1967) stated that Fabricius provided a diagnostic key for all generic names. Furthermore, some generic names were provided with descriptions or references to earlier published descriptions. Stafleu & Cowan (1976, pp. 810-811) concurred with Dandy's assessment that validation of Fabricius' generic names by means of author reference or by means of key are in full agreement with the *Code*. However, the nomenclatural validity of such names was questioned by Rauschert (1968) and Holub (1970).

In his detailed analysis of Fabricius' work, Holub (1970) concluded that not all of the generic names were validly published in that work. Regarding the generic names for which Fabricius either provided descriptions or references to earlier published descriptions, Holub considered them to be valid. Other names, appearing to be generic, but lacking validating descriptions, were treated by Holub to be uninomial designations of species (not true generic names; see *ICBN* Art. 20.4b, Ex. 10; Greuter 1988). Holub (pp. 84-86) divided all generic names (listed in Fabricius' work) into eight groups. He listed 120 invalid generic names (i.e., uninomial designations) within groups 1-6, and listed 32 valid generic names within groups 7 and 8. He treated the name *Laburnum* as a uninomial designation within group 3.

We concur with Holub's analysis, and treat *Laburnum* Fabr. as a unitary name. However, if some author were to use *Laburnum* Fabr. as a genus (prior to Medikus' 1787 work), then inadvertently, the unitary name would become a valid generic name. In the absence of such evidence, we accept *Laburnum* Medik. as a legitimate generic name.

Laburnum Medik., Vorles. Churpfalz. Phys.-Ocon. Ges. 2:363. 1787.

RANUNCULACEAE

Caltha palustris var. *palustris*, var. *radicans*, var. *flabellifolia*

Smit (1973, pp. 141-142) assigned the expression of *Caltha palustris* L., found mostly in Europe, Russia (?Cape Serdtze), Sakhalin, Kuril Islands, Japan, Alaska, and the Yukon territories of Canada, to var. *radicans* (Forst.) Beck. She cited *C. palustris* var. *flabellifolia* (Pursh) Torr. & Gray (based on *C. flabellifolia* Pursh) with its type from Pennsylvania, as a synonym. However, she (p. 143) did not include Pennsylvania within the range of var. *radicans*.

Gray (1895, p. 39) treated *Caltha palustris* var. *flabellifolia* as a synonym of the typical expression of *C. palustris*. Probably unaware of Beck's combination,

Gray proposed *C. palustris* var. *radicans* as a "new var." based upon *C. radicans*, and thus created an isonym.

Keener (Blumea 23:161. 1976.) stated that *Caltha palustris* var. *flabellifolia* is the correct name for the above complex, since the preceding varietal name has priority over *C. palustris* var. *radicans* at varietal rank (see below). Clapham (1987, p. 33), perhaps following Keener, assigned the British Isles' *C. palustris* [previously known as *C. palustris* var. *radicans* (Forst.) Beck] to var. *flabellifolia*. Akeroyd (in Tutin, et al. 1993, p. 254) merely mentioned the name *C. palustris* var. *radicans*, but did not mention the name *C. palustris* var. *flabellifolia*. Furthermore, he erred in attributing the name *C. palustris* var. *radicans* to (Forst.) Hook.

We believe that Gray was correct in his assessment of the type of *Caltha flabellifolia* belonging to *C. palustris* var. *palustris*. For the North American flora, we recognize two varieties within *C. palustris* as given below.

Caltha palustris L. var. *palustris*.

SY: *Caltha flabellifolia* Pursh, *Fl. Amer. Sept.* 390. 1814. *Caltha palustris* L. var. *flabellifolia* (Pursh) Torr. & Gray, *Fl. N. Amer.* 1:27. 1838.

Caltha palustris L. var. *radicans* (Forst.) Beck, *Verh. Zool. Bot. Gesellsch. Wien* 36:350. 1886. BASIONYM: *Caltha radicans* Forst., *Trans. Linn. Soc.* 8:324. 1807.

Ranunculus L.

Based on the belief that *Ranunculus triternatus* A. Gray (published in 1886) was a later homonym of *R. triternatus* Poir. (published in 1815), Nelson & Macbride (*Bot. Gaz.* 56:473. 1913.) proposed *R. reconditus* as a nom. nov. for Gray's taxon. When Benson (*Amer. J. Bot.* 23:170. 1936) reduced Gray's taxon as a variety under *R. glaberrimus* Hook., he believed that the epithet "triternatus" was not available for a transfer and therefore, chose the epithet "reconditus." However, eighteen years later, Benson (*Amer. Midl. Naturalist* 52:357. 1954.) realized that the name *R. triternatus* Poir. was invalidly published and that *R. triternatus* A. Gray was legitimate when published. This disposition rendered the name *R. reconditus* Nels. & Macbr. illegitimate (i.e., superfluous). Benson also realized that his earlier choice of the epithet "reconditus" at varietal rank was final; however, he failed to omit the parenthetical authorship for the var. *reconditus* (see *ICBN Art.* 72.2 Note 1). The correct authorship of var. *reconditus* is given below.

Ranunculus glaberrimus Hook. var. *reconditus* Benson, Amer. J. Bot. 23:170. 1936. BASIONYM: *Ranunculus reconditus* Nels. & Macbr., Bot. Gaz. 56:473. 1913, *nom. superfl.*

SY: *Ranunculus triternatus* A. Gray, Proc. Amer. Acad. Arts 21:370. 1886.

RHAMNACEAE

Frangula and *Rhamnus*

Ever since P. Miller (1768) proposed the genus *Frangula*, its disposition has varied. It was treated either as a subgenus or section of the genus *Rhamnus* L. (De Candolle 1825; Torrey & Gray 1838; Bentham & Hooker 1862; Weberbauer 1896; C. B. Wolf 1938; Johnston & Johnston 1978) or as a distinct genus (Martius 1861; V.I. Grubov 1949a, b; Tutin, *et al.* 1968; Weber 1990; Weber & Wittmann 1992).

Gray's treatment on the rank of *Frangula* varied. In Torrey & Gray (1838), *Frangula* was included within *Rhamnus*. However, in his independent works (1856-1867), Gray recognized *Frangula* as a distinct genus. But this position was reversed in Gray's 1889 work (which was revised by Watson & Coulter).

The traditional morphological separation of *Frangula* and *Rhamnus* has been as shown in Table 1 (based on our specimen observation and literature from the authors listed above).

Regarding the comparison of the above two genera, Johnston & Johnston (pp. 5-6) also provided a table summarizing the differences. They (p. 3) argued that *Frangula* "is exceedingly closely related to *Rhamnus*, more closely in fact than either group is to other taxa of the family." Furthermore, they stated that the recognition of a rank within a complex such as this "is not a scientific question, but one depending on tradition, usage and the practical consideration of the optimization of communication." They chose to treat *Frangula* as a subgenus within *Rhamnus*. These authors were critical of the European works on this complex remarking that "this tendency to recognize segregate genera seems to be fashionable in Europe."

Historically, the segregation or lumping of genera was an author's choice based primarily upon morphology. In modern literature, morphological differences are complemented with support from other areas, such as cytology, genetics, cladistics, phenetics, and protein and DNA analyses. For the segregation of *Frangula* from *Rhamnus*, Grubov (1949b) added the anatomical difference (*vide* Johnston & Johnston 1978): "the vessels have a different distribution in the xylem of *Frangula* than in that of the rest of *Rhamnus*."

In most of the recent North American floristic literature (excluding Weber 1990; Weber & Wittmann 1992), *Frangula* has been treated as a subgenus

Table 1. Comparison of *Rhamnus* and *Frangula*.

	<i>Frangula</i>	<i>Rhamnus</i>
Bud Scales	Absent	Present
Thorns	Absent	Present or absent
Leaf Pinnate Nerves	Almost straight	Often arcuate
Inflorescence	Umbels, fascicles, or flowers solitary	Racemes, fascicles, or flowers solitary
Flowers	Bisexual & mostly 5-merous	Mostly unisexual & 4-merous
Sepals	Mostly erect, fleshy keeled ventrally	Spreading, thin unkeeled
Petals	Well-developed, short-clawed	Usually small, often absent in pistillate
Anthers	Equalling filaments	Much shorter than filaments
Ovary	2 or 3-locular	2 or 4-locular
Style	Usually unbranched & not exerted	3- or 4-branched & exerted
Seeds	Smooth	Grooved outside
Raphe Position	Lateral	Apical
Cotyledons	Thick, straight	Thin, usually curved
Endosperm	Scanty	Abundant

within *Rhamnus*. However, we concur with Grubov (1949b), Tutin, *et al.* (1968), and Weber & Wittman (1992) in treating *Frangula* and *Rhamnus* as distinct genera. We assert that the differences summarized in the above table warrant the restoration of generic status to *Frangula*, which necessitates the following new combinations.

Frangula betulifolia (E. Greene) V. Grub. ssp. ***obovata*** (Kearney & Peebles) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus betulifolia* E. Greene var. *obovata* Kearney & Peebles, J. Wash. Acad. Sci. 29:486. 1939.

Frangula* × *blumeri (E. Greene) Kartesz & Gandhi, *comb. nov. (pro sp.)* [*betulifolia* × *californica*]. BASIONYM: *Rhamnus blumeri* E. Greene, Leaf. Bot. Obs. & Crit. 2:266. 1912.

Frangula californica (Eschsch.) A. Gray ssp. ***crassifolia*** (Jepson) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus californica* Eschsch. var. *crassifolia* Jepson, Fl. Pl. Calif. 615. 1925. SY: *Rhamnus californica* Eschsch. ssp. *crassifolia* (Jepson) C.B. Wolf, N. Amer. Sp. Rhamnus 68. 1938. *Rhamnus tomentella* Benth. ssp. *crassifolia* (Jepson) O. Sawyer, Madroño 40:65. 1993.

Frangula californica (Eschsch.) A. Gray ssp. **cuspidata** (E. Greene) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus cuspidata* E. Greene, Leafl. Bot. Obs. & Crit. 1:64. 1904. SY: *Rhamnus californica* Eschsch. ssp. *cuspidata* (E. Greene) C.B. Wolf, *N. Amer. Sp. Rhamnus* 72. 1938. *Rhamnus tomentella* Benth. ssp. *cuspidata* (E. Greene) O. Sawyer, *Madroño* 40:65. 1993.

Frangula californica (Eschsch.) A. Gray ssp. **occidentalis** (T.J. Howell) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus occidentalis* T.J. Howell, *Pittonia* 2:15. 1899. SY: *Rhamnus californica* Eschsch. ssp. *occidentalis* (T.J. Howell) C.B. Wolf, *N. Amer. Sp. Rhamnus* 66. 1938. *Rhamnus californica* Eschsch. var. *occidentalis* (T.J. Howell) Jepson, *Fl. Pl. Calif.* 615. 1925.

Frangula californica (Eschsch.) A. Gray ssp. **tomentella** (Benth.) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus tomentella* Benth., *Pl. Hartw.* 303. 1848. SY: *Rhamnus californica* Eschsch. ssp. *tomentella* (Benth.) C.B. Wolf, *N. Amer. Sp. Rhamnus* 70. 1938.

Frangula californica (Eschsch.) A. Gray ssp. **ursina** (E. Greene) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus ursina* E. Greene, Leafl. Bot. Obs. & Crit. 1:63. 1904. SY: *Rhamnus californica* Eschsch. ssp. *ursina* (E. Greene) C.B. Wolf, *N. Amer. Sp. Rhamnus* 74. 1938. *Rhamnus californica* Eschsch. var. *ursina* (E. Greene) McMinn, *Ill. Man. Calif. Shrubs* 329. 1939. *Rhamnus tomentella* Benth. ssp. *ursinus* (E. Greene) O. Sawyer, *Madroño* 40:65. 1993.

Frangula rubra (E. Greene) V. Grub. ssp. **modocensis** (C.B. Wolf) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus rubra* E. Greene ssp. *modocensis* C.B. Wolf, *N. Amer. Sp. Rhamnus* 89. 1938.

Frangula rubra (E. Greene) V. Grub. ssp. **nevadensis** (A. Nels.) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus nevadensis* A. Nels., *Proc. Biol. Soc. Wash.* 18:174. 1905. SY: *Rhamnus rubra* E. Greene ssp. *nevadensis* (A. Nels.) C.B. Wolf, *N. Amer. Sp. Rhamnus* 86. 1938.

Frangula rubra (E. Greene) V. Grub. ssp. **obtusissima** (E. Greene) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus obtusissima* E. Greene, Leafl. Bot. Obs. & Crit. 1:64. 1904. SY: *Rhamnus rubra* E. Greene ssp. *obtusissima* (E. Greene) C.B. Wolf, *N. Amer. Sp. Rhamnus* 88. 1938.

Frangula rubra (E. Greene) V. Grub. ssp. **yosemitana** (C.B. Wolf) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus rubra* E. Greene ssp. *yosemitana* C.B. Wolf, *N. Amer. Sp. Rhamnus* 90. 1938.

Frangula sphaerosperma (Sw.) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus sphaerosperma* Sw., *Prodr. Veg. Ind. Occ.* 50. 1788.

Frangula sphaerosperma (Sw.) Kartesz & Gandhi ssp. **longipes** (M.C. & L.A. Johnston) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus sphaerosperma* Sw. var. *longipes* M.C. & L.A. Johnston, *Fl. Neotrop. Monogr.* 20:62. 1978.

Rhamnus davurica and *R. citrifolia*

Hess & Stearn (1979) replaced the name *Rhamnus davurica* Pallas (published in 1776), a widespread species in southeastern Asia and introduced elsewhere, with their new combination *R. citrifolia* (Weston) Hess & Stearn, based on *Cornus citrifolia* Weston (published in 1770). These authors also made a new varietal combination in *R. citrifolia*: var. *nipponica* (Makino) Hess & Stearn, based on *R. davurica* var. *nipponica* Makino. Hess & Stearn's nomenclature was followed by Pratt (1980), and Gleason & Cronquist (1991, p. 342). Unfortunately, Hess & Stearn's combination is a later homonym of *R. citrifolia* Rusby (1907, p. 340), a South American taxon. These two identical names belong to two different genera: Weston's taxon is a *Rhamnus*, whereas Rusby's taxon is a *Frangula* (e.g., *F. citrifolia* (Rusby) V. Grub., *Trudy Bot. Inst. Akad. Nauk SSSR*, ser. 1. 8:276. 1949.).

Since Hess & Stearn's combination is illegitimate, the name *Rhamnus davurica* is reinstated here in place of *R. citrifolia* (Weston) Hess & Stearn (see the nomenclature given below), and a new subspecific combination is made here.

Rhamnus davurica Pallas, *Reise Prov. Russ.* 3:721. 1776 as 'davuricus'; Grubov, *Fl. URSS* 14:658, t. 36, f. 4. 1949. SY: *Rhamnus cathartica* L. var. *davurica* (Pallas) Maxim., *Mem. Acad. Sci. St. Petersburg.* 7(11):9. 1886. *Cornus citrifolia* Weston, *Univ. Bot.* 1:74. 1770. *Rhamnus citrifolia* (Weston) Hess & Stearn, *Taxon* 28:555. 1979, *non* Rusby, 1907.

Rhamnus davurica Pallas ssp. **nipponica** (Makino) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Rhamnus davurica* Pallas var. *nipponica* Makino, *Bot. Mag. Tokyo* 18:98. 1904. SY: *Rhamnus citrifolia* (Weston) Hess & Stearn var. *nipponica* (Makino) Hess & Stearn, *Taxon* 28:556. 1979.

Rhamnus lanceolata

Within *Rhamnus* species, we recognize subspecific (rather than varietal) rank, and therefore elevate *Rhamnus lanceolata* var. *glabrata* to subspecific rank as given below.

Rhamnus lanceolata Pursh ssp. **glabrata** (Gleason) Kartesz & Gandhi, *stat. nov.* BASIONYM: *Rhamnus lanceolata* Pursh var. *glabrata* Gleason, *Phytologia* 2:288. 1947.

ROSACEAE

The genus name *Chaenomeles* Lindl. was originally spelled *Choenomeles* by Lindley in 1821. In 1830, Bartling altered the name to *Chaenomeles*. This change in spelling was adopted by Lindley himself and most later authors. Wijandans (1990) proposed the conservation of the current usage of the spelling (i.e., *Chaenomeles*). However, based on *ICBN* Art. 73.3, we opted to use the original spelling, as did Zander's *Handwörterbuch*, ed. 13. 1984. The *ICBN* Committee for Spermatophyta (Taxon 42:877. 1993) recommended the conservation of the spelling *Chaenomeles*. The committee also stated that the change in spelling was allowed by Art. 73.1. We accept the committee's recommendation.

Chaenomeles Lindl., *Trans. Linn. Soc. London* 13:96. 1821. ("*Choenomeles*").

Prunus pumila var. *susquehanae*

In the protologue of *Prunus pumila* L. (the Sand-cherry), Gleason & Cronquist (1991, p. 260) included four varieties. For var. *cuneata* (Raf.) L.H. Bailey, they cited *P. susquehanae*, without authorship, as a synonym. However, Groh & Senn (1940, p. 332) used the name *P. pumila* L. var. *susquehanae* (hort. ex Willd.) Jaeg. and cited *P. pumila* var. *cuneata* (Raf.) Bailey as its synonym. We conclude that Groh & Senn were correct in their assessment as delineated below.

Prunus pumila L. var. *susquehanae* (hort. ex Willd.) Jaeg., *Ziergehoelze der Garten und Parkanlagen* 400. 1865. BASIONYM: *Prunus susquehanae* hort. ex Willd., *Enum. Pl. Hort. Berol.* 519. 1809. SY: *Prunus cuneata* Raf., *Ann. Nat.* 11: 1820. *Prunus pumila* L. var. *cuneata* (Raf.) L.H. Bailey, *Cycl. Amer. Hort.* 1451. 1901.

SOLANACEAE

Physalis ixocarpa, *P. philadelphica*

Regarding the occurrence of *Physalis ixocarpa* Brot. ex Hornem. in North America (north of México), Waterfall (1958, p. 160) gave its Canadian range as Ottawa and its US range as CA, DC, DE, MA, MD, NJ, NM, NY, OR, PA, TX, VA, VT, WA, and WV. He did not mention the species *P. philadelphica* Lam., which is found in México and Central America. However, in his treatment of *Physalis* in México, Central America, and West Indies, Waterfall (1967, p. 213) cited *P. ixocarpa* as a synonym of *P. philadelphica*. Furthermore, Waterfall recognized two varieties within *P. philadelphica*: var. *immaculata* and var. *parviflora*. Gleason & Cronquist (1991, p. 403) followed Waterfall's 1967 treatment and stated that *P. philadelphica* occasionally escaped from cultivation in the northeastern United States.

Based on her morphological and cytological analyses, Fernandes (1970, pp. 357-366) segregated *Physalis ixocarpa* from the *P. philadelphica* complex. She stated that *P. ixocarpa* has small flowers (calyx 4.00-5.25 mm long, 2.5-3.5 mm wide; corolla 5-10 mm in diameter; filaments 1.00-2.50(-2.75) mm long; anthers 1.25-1.75(-2.00) mm long; style 1.75-3.00 mm long), clavate stigma, and satellited chromosomes. The species *P. philadelphica* (mentioned as *P. ixocarpa* auct. non Brot. ex Hornem.) has large flowers (calyx ca. 8 mm long and 10 mm wide; corolla (10-)13-25(-30) mm in diameter; filaments ca. 5 mm long; anthers 3-4 (-5) mm long; style ca. 8 mm long), and lacks satellited chromosomes. Hudson (1983, p. 13; 1986, p. 417), who accepted Fernandes' conclusion, stated that *P. philadelphica* vars. *immaculata* and *parviflora* are self-compatible, and do not cross with *P. philadelphica* var. *philadelphica*, which is self-incompatible. He concluded that vars. *immaculata* and *parviflora* of *P. philadelphica* should be referred to *P. ixocarpa*.

Based on Fernandes' and Hudson's findings, we treat *Physalis ixocarpa* and *P. philadelphica* as two distinct species, and exclude *P. philadelphica* from North America, north of México. Furthermore, we transfer vars. *immaculata* and *parviflora* to *P. ixocarpa*, and make the two new combinations given below.

Physalis ixocarpa Brot. ex Hornem., *Suppl. Hort. Bot. Hafn.* 26. 1819.

Physalis ixocarpa* Brot. ex Hornem. var. *immaculata (Waterfall) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Physalis philadelphica* Lam. var. *immaculata* Waterfall, *Rhodora* 69:215. 1967.

Physalis ixocarpa* Brot. ex Hornem. var. *parviflora (Waterfall) Kartesz & Gandhi, *comb. nov.* BASIONYM: *Physalis philadelphica* Lam. var. *parviflora* Waterfall, *Rhodora* 69:215. 1967.

ACKNOWLEDGMENT

The authors thank Drs. Guy Nesom (TEX) and Larry E. Brown (Houston Community College) for their review of this article. The authors also thank Dr. Thomas Duncan (UC) for permitting the senior author to assess the specimens of *Frangula* and *Rhamnus*.

REFERENCES

- Bartram, W. 1791. *Travels Carolina*. James & Johnson, Philadelphia, Pennsylvania.
- Bentham, G. & J.D. Hooker. 1862-67. *Gen. Pl.* vol. 1. A. Black, London, Great Britain.
- Brummitt, R.K. 1993. *Vascular Plant Families and Genera*. Royal Botanic Gardens, Kew, Great Britain.
- Camp, W.H., et al. 1947. *International Rules of Botanical Nomenclature* (compiled from various sources). *Brittonia* 6:1-120.
- De Candolle, A.P. 1825. *Prodr.*, vol. 2. Treuttel & Wurtz, Paris, France.
- Clapham, A.R., et al. 1987. *Flora of British Isles*, ed. 3. Cambridge University Press, Cambridge, Great Britain.
- Dandy, J.E. 1967. Index of generic names of vascular plants 1753-74. *Regnum Veg.* vol. 51. Utrecht, The Netherlands.
- Fabricius, P.C. 1759. *Enum. Litteris Ioannis Drimbornii*, Helmstedt.
- Fernald, M.L. 1950. *Gray's Man. Bot.*, ed. 8. American Book Co., New York, New York.
- Fernandes, R.B. 1970. Sur l'identification d'une espee de *Physalis* sous-pontane au Portugal. *Bol. Soc. Brot.* 44:343-366.
- Gleason, H.A. & A. Cronquist. 1991. *Man. Vasc. Pl. NE US.*, ed. 2. The New York Botanical Garden, Bronx, New York.
- Gray, A. 1856. *Man. Bot.*, ed. 2. George P. Putnam & Co., New York, New York.
- . 1867. *Man. Bot.*, ed. 5. Ivison, Blakeman, Taylor, & Co., New York, New York.

- _____. 1889. *Man. Bot.*, ed. 6. American Book Company, New York, New York.
- _____. 1895. *Syn. Fl. N. Amer.*, vol. 1, pt. (1). New York, New York.
- Greuter, W. (Ch. Ed. Comm.). 1988. *International Code of Botanical Nomenclature*. Adopted by the Fourteenth International Botanical Congress, Berlin, Jul-Aug 1987. *Regnum Veg.* vol. 118. Utrecht, The Netherlands.
- _____, et al. 1989. Med - Checklist 4, Dicotyledons (Lauraceae - Rhamnaceae). Conservatoire et Jardin botaniques, Ville de Genève, Med-Checklist Trust of OPTIMA, Geneva, Switzerland.
- Groh, H. & H.A. Senn. 1940. *Prunus* in eastern Canada. *Canad. J. Res.* 18:318-346.
- Grubov, V.I. 1949a. Rhamnales. In: V.L. Komarov, *Fl. URSS* 14:635-674.
- _____. 1949b. *Trudy Bot. Inst. Akad. Nauk SSSR*, Ser. 1, Fl. Sist. Vyss. Rast.
- Hess, W.J. & W.T. Stearn. 1979. The identity of *Cornus citrifolia* Weston. *Taxon* 28:553-556.
- Holub, J. 1970. *Lamiastrum* versus *Galeobdolon* and comments on unitary designations in Fabricius's work "Enumeratio methodica plantarum horti medici helmstadiensis." *Folia Geobot. Phytotax.*, Praha 5:61-88.
- Hudson, W.D., Jr. 1983. The relationship of wild and domesticated tomato, *Physalis philadelphica*, Lamarck (Solanaceae). Ph.D. Dissert., Indiana University, Bloomington, Indiana.
- _____. 1986. Relationships of domesticated and wild *Physalis philadelphica*. pp. 416-432. In: W.G. D'Arcy (Ed.), *Solanaceae - Biology and Systematics*. Columbia University Press, New York, New York.
- Hutchinson, J. 1954. *Fl. W. Trop. Afr.*, vol. 1(1). Revised by R.W.J. Keay. Crown Agents, London, Great Britain.
- Jackson, B.D. 1895. *Index Kew*. The Clarendon Press, Oxford, Great Britain.
- Johnston, M.C. & L.A. Johnston. 1978. *Rhamnus*. *Fl. Neotropica Monogr.* No. 20. The New York Botanical Garden, Bronx, New York.

Kartesz, J.T. 1994. *A Synonymized Checklist of the Vascular Flora of United States, Canada & Greenland*, ed. 2. Timber Press, Inc., Portland, Oregon.

_____ & K.N. Gandhi. 1989. Nomenclatural notes for the North American flora - I. *Phytologia* 67:461-468.

_____. 1990a. Nomenclatural notes for the North American flora - II. *Phytologia* 68:421-427.

_____. 1990b. Nomenclatural notes for the North American flora - III. *Phytologia* 69:129-137.

_____. 1990c. Nomenclatural notes for the North American flora - IV. *Phytologia* 69:301-312.

_____. 1991a. Nomenclatural notes for the North American flora - V. *Phytologia* 70:194-208.

_____. 1991b. Nomenclatural notes for the North American flora - VI. *Phytologia* 71:58-65.

_____. 1991c. Nomenclatural notes for the North American flora - VII. *Phytologia* 71:87-100.

_____. 1991d. Nomenclatural notes for the North American flora - VIII. *Phytologia* 71:269-280.

_____. 1991e. Varietal autonyms of nontypical subspecies validly published combinations under today's Code. *Taxon* 40:308-310.

_____. 1992a. Nomenclatural notes for the North American flora - IX. *Phytologia* 72:17-30.

_____. 1992b. Nomenclatural notes for the North American flora - X. *Phytologia* 72:80-92.

_____. 1992c. Nomenclatural notes for the North American flora - XI. *Phytologia* 73:124-136.

_____. 1992c. Nomenclatural notes for the North American flora - XII. *Phytologia* 74:43-55.

Kotschy, K.G.T. & J.J. Peyritsch. Jul 1867. *Pl. Tinn.* Vindobonae (Wien), Austria.

- Leadlay, E.A. & V.H. Heywood. 1990. The biology and systematics of the genus *Coincya* Porta & Rigo ex Rouy (Cruciferae). *J. Linn. Soc. Bot.* 102:313-398.
- Linnaeus, C. 1753. *Sp. Pl. Impensis Laurenti Salvii*, Stockholm, Sweden.
- Martius, C.F.P. 1861. *Fl. Bras.*, vol. 11(1). Leipzig, Germany. Facsimile ed. issued in 1965 by J. Cramer, Weinheim, Germany.
- Meeuse, A.D.J. 1962. The Cucurbitaceae of South Africa. *Bothalia* 8:1-111.
- Merrill, E.D. 1945. In defence of the validity of William Bartram's binomials. *Bartonia* 23:10-35.
- Miller, P. 1768. *Gard. Dict.*, ed. 8. London, Great Britain.
- Pratt, J.D. 1980. New England Note: a naturalized population of *Rhamnus citrifolia* in Connecticut. *Rhodora* 82:523-524.
- Rauschert, S. 1968. Zur frage der gattungsnamen bei Fabricius. *Taxon* 17:153-156.
- Rickett, H.W. 1944. Legitimacy of names in Bartram's "Travels." *Rhodora* 46:389-391.
- Rollins, R.C. 1993. *The Cruciferae of Continental North America*. Stanford University Press, Stanford, California.
- Rusby, H.H. 1905. An enumeration of the plants collected in Bolivia by Miguel Bang, part. 4. *Bull. New York Bot. Garden* 4(14):309-470.
- Smit, P.G. 1973. A revision of *Caltha* (Ranunculaceae). *Blumea* 21:119-150.
- Stafleu, F.A. & Cowan. 1976. *Taxonomic Literature*, ed. 2, vol. 1:A-G. *Regnum Veg.* vol. 94. Utrecht, The Netherlands.
- _____. 1979. *Taxonomic Literature*, ed. 2, vol. 2: H-Le. *Regnum Veg.* vol. 98. Utrecht, The Netherlands.
- Torrey, J. & A. Gray. 1838. *Fl. N. Amer.*, vol. 1(2). G. & C. Carvill & Co., New York, New York.
- Tutin, T.G., et al. (Eds.). 1968. *Fl. Europaea*, vol. 2. Cambridge University Press, Cambridge, Great Britain.
- _____ et al. (Eds.). 1993. *Fl. Europaea*, vol. 1, ed. 2. Cambridge University Press, Cambridge, Great Britain.

- Voss, E.G. 1985. Michigan Flora, Part 2: Dicots (Saururaceae - Cornaceae). Cranbrook Institute of Science Bull. 59.
- Waterfall, U.T. 1958. A taxonomic study of the genus *Physalis* in North America north of Mexico. *Rhodora* 60:107-114, 128-142, 152-173.
- . 1967. *Physalis* in Mexico, Central America, and the West Indies. *Rhodora* 69:88-120; 203-239; 319-329.
- Weber, W.A. 1990. *Colorado Flora: Eastern Slope*. University Press of Colorado, Niwot, Colorado.
- & R.C. Wittmann. 1992. *Catalog of the Colorado Flora: A Biodiversity Baseline*. University Press of Colorado, Niwot, Colorado.
- Weberbauer, A. 1895-96. Rhamnaceae. Pp. 393-427. In: Engler, A. & K. Prantl (Eds.), *Die Natürlichen Pflanzenfamilien*, Teil. vol. 3, Abteilung vol. 5, Lieferung part 128, 136. Verlag von Wilhelm Engelmann, Leipzig, Germany.
- Welman, W.G. 1987. Convolvulaceae to Asteraceae. Pp. 163-253. In: G.E. Gibbs Russell, *et al.* (eds.), *List of Species of Southern African Plants*, ed. 2, pt. 2. Botanical Research Institute, Dept. of Agriculture and Water Supply, Pretoria, South Africa.
- Wiersema, J.H., *et al.* 1990. Legume (Fabaceae) Nomenclature in the USDA Germplasm System. USDA, Agr. Res. Serv. Tech. Bull. No. 1757.
- Wijnands, D.O. 1990. Proposal to conserve the spelling of 3336a *Chaenomeles* Lindley (Rosaceae). *Taxon* 39:534-535.
- Wilbur, R. 1971. A reconsideration of Bartram's binomials. *J. Elisha Mitchell Soc.* 87:56-73.
- Wolf, C.B. 1938. *N. Amer. Sp. Rhamnus*. Rancho Santa Ana Bot. Gard. Monogr. Bot. Ser. No. 1.

**A NEW LICHEN SPECIES, *NIEBLA CEDROSENSIS* (RAMALINACEAE), IS
DESCRIBED FROM BAJA CALIFORNIA, MEXICO**

J.E. Marsh & T.H. Nash, III

Department of Botany, Arizona State University, Tempe, Arizona 85287-1601
U.S.A.

ABSTRACT

A new species, *Niebla cedrosensis* Marsh & Nash, *sp. nov.*, is described from Baja California, México. This saxicolous species has pendant terete branches, black pycnidia, terminal to subterminal apothecia and adglutinated hyphae in the medulla. At its center of distribution on Isla Cedros, the thallus has a distinctive pale yellow-green color due to cortex abrasion.

KEY WORDS: lichen, Ramalinaceae, Baja California

Niebla cedrosensis Marsh & Nash, *sp. nov.* (Figure 1). TYPE: MEXICO. Baja California. Isla Cedros at southwest end, NE of Wayle, 28°07' N 115°19' W: on siliceous pebble on steep hillside, N aspect, 150 m elevation. March 22, 1994. *Marsh 7460* (ASU). Isotype (1): (ASU).

Description: Thallus saxicolus, pendulus usque ad 6.0 cm longus, luteo-viridis pallido; rami teretes ad 1.5 mm lati; medulla densa cum hyphae adglutinae continuae.

Thallus fruticose, saxicolous, pendant to 4.5 cm (-10.5 cm) long, 1.5 mm wide, stiff terete branches from a thick narrowly attached holdfast. Cortex smooth/foveate/reticulate, sometimes transversely cracked. Medulla solid, of adglutinated hyphae without distinctive chondroid strands, as is the case in the *Niebla homalea* group. Pycnidia black, on upper two-thirds of branch. Apothecia terminal to subterminal, to 4.5 mm diameter, multiple, disc white, pruinose; spores 8, straight to slightly curved, 2-celled, hyaline, 10.0-14.0 × 3.0-4.0 μm.




Figure 1. *Niebla cedrosensis*, (Type in ASU). (Scale=mm).




Figure 2. Distribution of *Niebla cedrosensis*.

Chemistry: \pm usnic, 16- α -hydroxykurane, zeorin, unknown triterpene, \pm salazinic acid

Habitat and Distribution: *Niebla cedrosensis* is saxicolous on sides and overhangs of boulders and rock outcrops. It is restricted to Cedros Island and the tip of the Vizcaino Peninsula (Figure 2) at 40-500 m elevation. Wind and salt and/or sand have abraded the thalli of this species so that it has a distinctive pale whitish yellow-green appearance and hence stands out from all other *Niebla/Ramalina* species. One yellowish-green non-abraded collection was found from Bahía Tortuga at the north-western tip of the Vizcaino Peninsula (13 miles southeast).

Representative Specimens Examined: MEXICO. Baja California: Isla Cedros, north end, Marsh 7194, 7283, 7294, 7295, 7306, 7309, 7339, 7342, 7344 (ASU); Nash 5 collections s.n., southwest end, Marsh 7418, 7446, 7458, 7523 (ASU). Bahía Tortuga, Marsh 4279 (ASU).

Remarks: Since the distribution of *Niebla cedrosensis* is restricted to Cedros Island and the tip of the Vizcaino Peninsula, this species should not be confused in the field. Other saxicolous *Niebla* species whose distributions overlap that of *N. cedrosensis* are *N. josecuervo* (Rundel & Bowler) Rundel & Bowler having only salazinic acid (Rundel *et al.* 1972) and flat branches; this species has two morphologies in this area, 1) long thin spirally twisted branches, 2) broad fan-like branches. Another species being described separately with similar chemistry is the same size as *N. cedrosensis*, but has sordid green branches, usually with black necrotic tissue at the base and lower portion of the branch and lacks adglutinated medullary hyphae.

ACKNOWLEDGMENTS

Mr. Jon Rebman and Dr. José Delgadillo, Universidad Autónoma de Baja California, Ensenada, Baja California, México who made initial arrangements for the field trip to Cedros Island and Biol. René Luis Aguilar Vásquez, Oficina Federal de Pesca and Sr. Rafael Muñoz Gallardo, Soc. Coop. Pescadores Nacionales de Abulon, S.C.L., Isla de Cedros, B.C., México for accommodation and transportation on the island. This research was supported by NSF grant #BSR9201111 to T.H. Nash III.

LITERATURE CITED

- Rundel, P.W., P.A. Bowler, & T.W. Mulroy. 1972. A fog-induced lichen community in northwestern Baja California, with two new species of *Desmazieria*. *The Bryologist* 75:501-508.

WEED FLORA OF KHAMMAM DISTRICT IN ANDHRA PRADESH, INDIA

G. Jaya Babu, V. Pushpalatha, & K.C. Naidu

Department of Botany, Andhra University, Waltair, INDIA

ABSTRACT

A summary of the weed flora of Khammam District is provided.

KEY WORDS: Floristics, invasive species, India

INTRODUCTION

Khammam District in Andhra Pradesh lies between 16° 45' and 18° 35' north latitude and between 79° 47' and 80° 47' east longitude. The district ranks sixth in the state with respect to area, covering 15,921 square kilometers, and is very rich in vegetation—particularly the ground flora. Khammam District has not been well explored floristically. A few botanists made random collections during brief periods over the past 90 years (Partridge 1911; Sayeedud-Din 1935-1941; Saxena 1944-1946; Khan 1953). Therefore, the present work is undertaken to summarize detailed observations on the wealth of weed species and floristic components.

ENUMERATION OF SPECIES

Identifications of plant material collected were made with help of *Flora of Madras Presidency* and *Flora of British India*, and later authenticated by the botanical Survey of India.

ACANTHACEAE: *Andrographis echiioides* L.; *Andrographis paniculata* (Burn. f.) Nees; *Blepharis asperrima* Nees; *Barleria strigosa* Wall.; *Dipterocanthus prostratus* (Poir.) Nees; *Justicia glauca* Rottl.; *Justicia procumbens* L.; *Justicia simplex* D. Don. *Ruellia tuberosa* L.; *Rungia pectinata* (L.) Nees; *Rungia repens* Nees.

- AIZOACEAE: *Giseckia pharmacoides* L. *Glinus lotoides* L.; *Glinus oppositifolius* (L.) A. DC.; *Mollugo nudicaulis* Lamk.; *Mollugo pentaphylla* L.; *Trianthema portulacastrum* L.
- AMARANTHACEAE: *Achyranthes aspera* L.; *Aerva lanata* (L.) Juss.; *Alla-
mania nodiflora* (L.) R.Br.; *Alternanthera sessilis* (L.) R.Br. *Amaranthus
spinosus* L.; *Digera muricata* (L.) Beitr.; *Gomphrena decumbens* Jacq.
- APOCYNACEAE: *Catharanthus pusillus* (Murr.) G. Don; *Catharanthus roseus*
(L.) G. Don.
- ASCLEPIADACEAE: *Calotropis gigantea* (L.) R.Br.; *Pergularia aemia* (Forsk.)
Chiov.; *Tylophora indica* (Burn. f.) Merrill.
- ASTERACEAE: *Acanthospermum hispidum* DC.; *Ageratum conyzoides* L.;
Caesulia azillaris Roxb.; *Eclipta alba* L.; *Emilia sonchifolia* (L.) DC.;
Parthenium hysterophorus L. *Sonchus oleraceus* L.; *Sphaeranthus indi-
cus* L.; *Tridax procumbens* L.; *Vernonia cinerea* (L.) Less.; *Xanthium
strumarium* L.
- BORAGINACEAE: *Coldenia procumbens* L.; *Heliotropium indicum* L.; *He-
liotropium scabrum* Retz.; *Trichodesma indicum* (L.) R.Br.
- CAESALPINACEAE: *Cassia absus* L. *Cassia occidentalis* L.; *Cassia sophora*
L.; *Cassia tora* L.
- CAPPARIDACEAE: *Capparis zeylanica* L. *Cleome aspera* Koenig.; *Cleome
monophylla* L.; *Cleome viscosa* L.; *Gynandropsis pentaphylla* (L.) DC.
- COMMELINACEAE: *Commelina attenuata* Koenig; *Commelina bengalen-
sis* L.; *Commelina erecta* L.; *Commelina longifolia* Lamk.; *Cyanotis az-
illaris* Roem. & Schult.
- CONVOLVULACEAE: *Ipomoea carnea* Jacq.; *Ipomoea hederifolia* L.; *Ipo-
moea nil* (L.) Roth.; *Ipomoea pestigradis* L.; *Merremia emarginata* (Burn.
f.) Hall; *Merremia tridentata* (L.) Hall.
- CUCURBITACEAE: *Bryonopsis laciniosa* (L.) Naud.; *Ctenolepis garcini*
Naud. *Cucumis trigonus* Roxb.; *Mukia scabrella* Arn.
- CYPERACEAE: *Cyperus alopecuroides* Roth.; *Cyperus cephalotus* Vahl.;
Cyperus compactus Retz.; *Cyperus difformis* L.; *Cyperus distans* L.;
Cyperus iria L.; *Cyperus platystylis* R.Br.; *Cyperus triceps* (Rottb.) Endl.;
Fimbristylis dichotoma (L.) Vahl.; *Fimbristylis ovata* Burn. f.

EUPHORBIACEAE: *Acalypha indica* L.; *Chrozophora rottleri* (Geiss.) Juss.; *Croton bonplandianum* Baill.; *Euphorbia hirta* L.; *Micrococca mercurialis* (L.) Benth.; *Phyllanthus amarus* Schum. & Thorn.; *Phyllanthus debilis* Klein.; *Phyllanthus maderaspatensis* L.; *Phyllanthus virgatus* Forst.; *Sebastiania chamaelea* (L.) Muell.

FABACEAE: *Alysicarpus monilifer* (L.) DC.; *Atylosia rugosa* W. & A.; *Atylosia scarabaeoides* (L.) Benth.; *Crotalaria linifolia* L.; *Crotalaria nana* Burn.; *Crotalaria prostrata* Rottl.; *Crotalaria verrucosa* L.; *Desmodium gangetium* (L.) DC.; *Desmodium triflorum* (L.) DC.; *Rhynchosia minima* (L.) DC. *Rhynchosia suaveolens* (L. f.) DC.; *Tephrosia hirta* Ham.; *Tephrosia procumbens* Buch-Ham.; *Tephrosia purpurea* L.; *Zornia diphylla* Pers.

LAMIACEAE: *Acrocephalus indicus* (Burn. f.) O. Ktze.; *Anisochilus carnosus* (L.) Wall.; *Anisomeles indica* (L.) O. Ktze.; *Hyptis suaveolens* (L.) Poir.; *Leucas aspera* (Willd.) Spr. *Ocimum americanum* L.; *Ocimum basilicum* L.

MALVACEAE: *Pavonia zeylanica* Con. *Sida acuta* Burn. f.; *Sida cordifolia* L.; *Sida rhombifolia* L.; *Sida veronicifolia* Lamk.; *Urena lobata* L.

PAPAVERACEAE: *Argemone mexicana* L.

PEDALIACEAE: *Martynia annua* L. *Petalium murex* L.

POACEAE: *Apluda mutica* L.; *Aristida funiculata* Trin. & Rupr.; *Brachiaria distachya* L.; *Cynodon dactylon* (L.) Pers.; *Dactyloctenium aegyptium* (L.) Beauv. *Eragrostiella bifaria* (Vahl.) Bor.; *Eragrostis tenella* (L.) Beauv.; *Eragrostis unioides* Nees; *Panicum repens* L.; *Perotis indica* (L.) O. Ktze.; *Themeda triandra* Forsk.; *Urochloa panicoides* Beauv.

PORTULACACEAE: *Portulaca oleracea* L.; *Portulaca quadrifida* L.

RUBIACEAE: *Borreria articularis* (L. f.) Williams.; *Borreria stricta* (L.) K. Schum.; *Galium asperifolium* Wall. *Knoxia sumatrensis* (Retz.) DC.; *Oldenlandia corymbosa* L.; *Oldenlandia herbacea* (L.) Roxb.; *Oldenlandia umbellata* L.

SAPINDACEAE: *Cardiospermum halicacabum* L.; *Dodonaea viscosa* (L.) Jacq.

SCROPHULARIACEAE: *Bacopa monnieri* (L.) Pennell; *Lindernia crustacea* (L.) F. Muller; *Scoparia dulcis* L.; *Stemodia viscosa* Roxb.; *Striga densiflora* Benth.

SOLANACEAE: *Physalis minima* L. *Solanum nigrum* L.; *Solanum surattense* Burn.; *Solanum trilobatum* L.

STERCULIACEAE: *Buettneria herbacea* Roxb.; *Helicteres isora* L.; *Melhania cannabina* Wight. *Waltheria americana* L.

TILIACEAE: *Corchorus aestuans* L. *Triumfetta pentandra* A. Rich.; *Triumfetta rhomboidea* Jacq.

VERBENACEAE: *Lantana camara* L.; *Lippia nodiflora* Mich.

ZYGOPHYLLACEAE: *Tribulus terrestris* L.

REFERENCES

- Gamble, J.S. & C.E.L. Fischer. 1915-1936. *Flora of the Presidency of Madras*. 3 vols. London. (reprinted in 1957). Botanical Survey of India, Calcutta, India.
- Hooker, J.D. 1872-1897. *The Flora of British India* 7 vols. London (reprinted in 1973). Bishen Singh Mahendrapal Singh, Dehra Dun and Periodical Experts, New Delhi, India
- Khan, M.S. 1953. *Forest Flora of Hyderabad State*. Government Press, Hyderabad, India.
- Partridge, E.A. 1911. *Forest Flora of H.E.H. the Nizam's Dominions, Hyderabad Deccan*. A. Venugopal Pillai & Sons, Hyderabad, India.
- Sayedud-Din, M. 1935. Some of the common flowering plants of Hyderabad, Their distribution, economic and medicinal importance. *J. Asiatic Soc. Bengal Sci.*, Calcutta.
- Sayedud-Din, M. 1938. Further contribution to some of the common flowering plants of the Hyderabad State, Their distribution and economic importance of dicots. *J. Bombay Nat. Hist. Soc.*
- Sayedud-Din, M. 1941. Additions to our knowledge of the flowering plants of H.E.H. the Nizam's Dominions, Hyderabad Deccan. *J. Bombay Nat. Hist. Soc.*

A NEW SPECIES OF *CASTILLEJA* (SCROPHULARIACEAE) RELATED TO *C. ASPERA* FROM SOUTHEASTERN DURANGO, MEXICO

Guy L. Nesom

Department of Botany, University of Texas, Austin, Texas 78713 U.S.A.

ABSTRACT

Castilleja gonzalezii, *sp. nov.*, is described from southeastern Durango, México. It is most closely related to three other species from the Sierra Madre Occidental with evenly divided, acute-lobed calyces, *C. aspera*, *C. chlorosceptron*, and *C. durangensis*. The new species differs from all three of these particularly in its densely stipitate-glandular vestiture.

KEY WORDS: *Castilleja*, Scrophulariaceae, Durango, México

Among recently incoming material at TEX is a specimen of *Castilleja* from southeastern Durango, México, that apparently represents a remarkably distinct, undescribed species.

Castilleja gonzalezii Nesom, *sp. nov.* TYPE: MEXICO. Durango: Mpio. Mezquitil, 33 km de la Guajolota por el camino a Platanitos; bosque abierto de pino-encino, 2220 m, 16 Mar 1985, *M. González et al. 1699* (HOLOTYPE: TEX!; Isotype: CIIDIR).

Castillejae durangensi Nesom similis sed differt vestimento dense stipitati-glanduloso, calycibus longioribus, et bracteis floralibus angustioribus.

Plants apparently perennial. Stems erect, 35 cm tall, broken off at the base but apparently several-stemmed from the base, probably taprooted, simple or with a single branch; stems, leaves, floral bracts, and calyces densely stipitate-glandular with hairs mostly 0.1-0.5 mm long, the glandular trichome apex orangish, eglandular hairs absent or rarely a few present along the leaf margins. Leaves erect-ascending, sometimes slightly recurved, linear-lanceolate, entire,

strongly 3-veined, the lowermost subclasping, non-clasping above, mostly 25-40 mm long, slightly shorter upwards, 2-4 mm wide (at midpoint), often folded at the middle. Mature inflorescence and infructescence ca. 8-10 cm long; floral bracts entire, 3-veined, 2.0-2.5 cm long, lanceolate, abruptly tapered from the base, 1.5-2.5 mm wide at midpoint, reddish or with a red apex, the apex slightly dilated. Calyces pink to red, 22-30 mm long, tubular (in fruit basally inflated and chartaceous, immediately constricted above the fruit), the primary lobes 10-14 mm long, nearly equal in length or the abaxial sinus slightly deeper, the secondary lobes 1-2 mm long, narrowly triangular. Corollas 24-37 mm long, the lower lip of 3, thick, green teeth ca. 1 mm long, the galea 12-17 mm long, sparsely glandular-pilose dorsally, 43-46% as long as the corolla, exerted 0-6 mm from the calyx; stigma bilobed, not or barely exerted.

Castilleja gonzalezii, known only from the type collection, is named for its collector, Martha González-Elizondo. She and her sister, Socorro González-Elizondo, have produced a detailed study of the vegetation of La Reserva de la Biosfera "La Michilia" in generally the same area of southeastern Durango where the new species was collected.

The new species is a member of *Castilleja* sect. *Euchroma* Benth. (sensu Eastwood 1909), which is characterized by an equally divided calyx and by the corolla tube and galea about equal in length. Within sect. *Euchroma*, *C. gonzalezii* is part of a small subgroup of four species with distinctly triangular, sharply acute, calyx lobes. All four of these species are restricted to the Sierra Madre Occidental of México: *C. aspera* Eastwood and *C. durangensis* Nesom (Nesom 1992a), and *C. chlorosceptron* Nesom (Nesom 1992b). *Castilleja gonzalezii* is most similar to *C. durangensis* and *C. chlorosceptron* in its linear, densely arranged leaves, but it differs from all three of its close relatives in its densely stipitate-glandular vestiture, the others being essentially eglandular. *Castilleja chlorosceptron* is diffusely rhizomatous and produces glabrous leaves and very small flowers (calyces 12-13 mm long, corollas 15-16 mm long); *C. durangensis* is taprooted and produces densely hispidulous leaves, small flowers (calyces 18-22 mm long, corollas 21-25 mm long), and floral bracts distinctly broadened at the apex.

ACKNOWLEDGMENTS

I thank B.L. Turner and Piero Delprete for their comments on the manuscript.

LITERATURE CITED

- Eastwood, A. 1909. Synopsis of the Mexican and Central American species of *Castilleja*. Proc. Amer. Acad. Arts 44:563-591.

- González-Elizondo, S. & M. González-Elizondo. 1993. Vegetación de la Reserva de la Biosfera "La Michilia," Durango, México. *Acta Bot. Mex.* 22:1-104.
- Nesom, G.L. 1992a. New species and taxonomic evaluations of Mexican *Castilleja* (Scrophulariaceae). *Phytologia* 72:231-252.
- . 1992b. A new species of *Castilleja* (Scrophulariaceae) from Durango, México. *Phytologia* 73:137-139.

TWO NEW SPECIES OF *SISYRINCHIUM* SUBG. *ECHTHRONEMA*
(IRIDACEAE) FROM NUEVO LEON, MEXICO

Guy L. Nesom

Department of Botany, University of Texas, Austin, Texas 78713 U.S.A.

ABSTRACT

Two new species of *Sisyrinchium* subg. *Echthronema* are described from Nuevo León, México. *Sisyrinchium hintoniorum* is a high elevation endemic most closely related to the more widespread *S. schaffneri*; *S. microbracteatum* apparently is restricted to gypsum outcrops and is most similar to *S. tenuifolium*.

KEY WORDS: *Sisyrinchium*, Iridaceae, México

Two remarkably distinct species of *Sisyrinchium* have been brought to attention in recent collections by the Hinton family from the state of Nuevo León in northeastern México. Flowers of both species are yellow and produce spreading, free staminal filaments, marking them as members of subg. *Echthronema* (Herbert) Goldblatt. Recent publications regarding the taxonomy of Mexican and Central American *Sisyrinchium* were summarized by Nesom & Hernández (1992) in the introduction to the description of a new species of subg. *Sisyrinchium* from Nuevo León.

Sisyrinchium hintoniorum Nesom, *sp. nov.* TYPE: MEXICO. Nuevo León, Mpio. Zaragoza, Cerro El Viejo, pine woods, 3125 m, 10 May 1992 [flowering], Hinton *et al.* 22440 (HOLOTYPE: TEX!).

Sisyrinchio schaffneri S. Wats. similis sed differt floribus numerosioribus, corollis multo majoribus, bracteis spathae longioribus, et seminibus majoribus sine depressione distincta.

Perennials from a short (5-10 mm), densely fibrous-rooted rhizome or caudex, the roots slightly thickened but not distinctly swollen, herbage glabrous. Stems 4-5 per plant, erect, 28-45 cm tall, flattened and 2-winged, 1.5-2.0 mm

wide, unbranched, bearing a single spathe at the apex. Leaves linear, linear-lanceolate apically, 1.5-4.0 mm wide, 2.5-3.5 cm long. Spathes solitary, 6-8-flowered, compressed, narrowly lanceolate-elliptic in outline, 3-4 mm wide, the spathe bracts subequal to unequal in length, the outer 44-55 mm long and 35-50% longer than the inner, the inner 30-35 mm long, the outer with basally connate margins 4-9 mm above the base. Tepals yellow, broadly ovate-lanceolate, 18-20 mm long; filaments united ca. 1.5 mm, free ca. 4 mm; anthers 5 mm long; ovaries obpyriform, glabrous. Capsules obovate-oblong in outline, 7-9 mm long, 4-5 mm wide, apically truncate, 3-angled, the surface often distinctly shiny; pedicels strongly nodding in fruit, ca. 20 mm long. Seeds black, distinctly foveolate, 1.5-2.0 mm in diameter, spheroid to slightly angular, without a distinct depression or the depression very shallow and ca. 1/3 of seed width.

Additional collection examined: MEXICO. Nuevo León: Mpio. Aramberri, Cerro El Viejo, dense "ayarin" forest, 3225 m, 20 Nov 1993 [fruiting only], Hinton *et al.* 23975 (TEX).

Within subg. *Echthronema*, the new species is similar and clearly closely related to three other species primarily centered in northern México: (1) *Sisyrinchium schaffneri* S. Wats., known from Arizona, Chihuahua, Coahuila, Nuevo León, Tamaulipas, San Luis Potosí (the type), Edo. México, Puebla, Michoacán, Jalisco, and Aguascalientes; (2) *S. cernuum* (Bickn.) Kearney, known from Arizona, Texas, Chihuahua, Coahuila, Nuevo León, Tamaulipas, Veracruz, Hidalgo, Querétaro, Puebla, Edo. México, Guanajuato, Aguascalientes, Jalisco, Sinaloa, and Durango; and (3) *S. longipes* (Bickn.) Kearney & Peebles, known from Arizona, Sonora, and Chihuahua. Each of these species produces unbranched stems with a single terminal spathe, spathe bracts of subequal to unequal length, and distinctly unthickened fibrous roots. These species can be distinguished by the contrasts in the following key.

1. Pedicels erect or ascending; tepals 8-12 mm long. *S. longipes*
1. Pedicels distinctly arcuate-nodding; tepals 3-5, 10-13, or 18-20 mm long.
..... (2)
 2. Tepals 3-5 mm long. *S. cernuum*
 2. Tepals 10-20 mm long. (3)
3. Tepals 10-13 mm long; flowers 2-4(-6) per spathe, the longer spathe bract 22-30(-35) mm long, shorter bract 20-26 mm long; seeds 1.2-1.5 mm wide, depressed-spherical, with a deep and sharply defined depression ca. 2/3 the seed width. *S. schaffneri*
3. Tepals 18-20 mm long; flowers 6-8 per spathe, the longer spathe bract 44-55 mm long, shorter bract 30-35 mm long; seeds 1.5-2.0 mm wide, rounded

to somewhat angular, without a distinct depression or the depression 1/3 the seed width and very shallow. *S. hintoniorum*

Of these, only *Sisyrinchium hintoniorum* and *S. schaffneri* occur in Nuevo León. The latter occurs at 1550-2380 meters on Cerro Viejo, apparently not reaching the high elevation of that mountain where *S. hintoniorum* grows, but it seems clear that *S. hintoniorum* cannot be considered merely a large, ecological variant of *S. schaffneri*. *Sisyrinchium schaffneri* is found at and near the top of two other high peaks in the area, Peña Nevada and Cerro Potosí, each of which is near 3300 meters.

***Sisyrinchium microbracteatum* Nesom, sp. nov.** TYPE: MEXICO. Nuevo León, Mpio. Aramberri, Aramberri to El Salitre, gypsum hillside, 1325 m, 26 Oct 1993, *Hinton et al. 29797* (HOLOTYPE: TEX!; Isotypes: CAS!, ENCB, GH!, MEXU!, NY!, US!).

Sisyrinchio tenuifolio Willd. similis sed differt caulibus elatioribus, bracteis spathae libris ad basim, bracteis pusillis ad nodos ac internodos superos, ovariis glabris, et capsulis minoribus laevibus.

Perennials without an evident caudex, with a fascicle of fleshy-thickened fibrous roots ca. 4-8 cm long, 2-4 mm wide, tapered slightly at the tips; herbage glabrous. Stems mostly 1-3 per plant, wiry-thin, terete, erect, 25-40 cm tall, (2-)3-5 branched in the upper half, each branch node with a folded bract (similar to outer spathe bract) 5-15 mm long, diminishing in size upward, a small peduncular bract commonly present 10-35 mm immediately below the spathe. Leaves linear, not folded basally, 1.0-2.5 mm wide, the basal 15-30 cm long, leaving a mass of persistent, fibrous bases mostly 2-4 cm long, cauline leaves shorter and 1-3 in number. Spathes solitary, 4-9-flowered, slightly compressed, elliptic to elliptic-lanceolate in outline, 2-4 mm wide, the spathe bracts equal, 10-14 mm long, or the outer 1/2-2/3 the length of the inner, the margins usually free to the very base but sometimes connate 0.3-0.6 mm, brownish-green, often purplish along the edges, the inner bract with distinctly hyaline margins. Tepals yellow, broadly obovate-oblancheolate, 9-12 mm long; filaments and style united into a column 1.0-1.5 mm long, the filaments spreading and free 2.0-2.5 mm, style branches 2.0-2.5 mm long, anthers ca. 2 mm long, strongly curved; ovaries spherical, glabrous. Capsules (mature) slightly oblong in outline, 3.5-5.5 mm long, 2.5-3.5 mm wide; pedicels erect, extending 0-4 mm above the spathe apex. Seeds black, very shallowly foveolate-reticulate, ca. 1 mm in diameter, spheroid, slightly depressed, with a deep, distinctly margined depression ca. 2/3 the width of the seed.

Additional collections examined: MEXICO. Nuevo León: Mpio. Galeana, near Río de San José, oak and pine woods, 1720 m, 2 Oct 1991, *Hinton et al.* 21570 (TEX); Mpio. Zaragoza, near Zaragoza, gypsum hillside, 1365 m, 28 Jul 1993, *Hinton et al.* 23104 (TEX); Mpio. Zaragoza, Cerro Viejo, gypsum hillside, 1350 m, 23 Sep 1993, *Hinton et al.* 23515 (TEX); Mpio. Aramberri, Aramberri to El Salitre, gypsum hillside, 1325 m, 26 Oct 1993, *Hinton et al.* 23736 (TEX).

Within *Sisyrinchium* subg. *Echthronema*, *S. microbracteatum* is similar and apparently most closely related to *S. tenuifolium* Willd., a widespread species in México. Both species produce branched, wingless, nearly terete stems, narrow leaves, a fascicle of fleshy-thickened fibrous roots, subequal spathe bracts, and at least some forms of *S. tenuifolium* have persistent, fibrous leaf bases. The new species differs from *S. tenuifolium* in its generally taller stems (vs. stems mostly 15-25 cm high), spathe bracts free to the base (vs. margins connate to 1.5-3.0 mm above the base), numerous small bracts at the branch nodes and on the peduncles (vs. lanceolate nodal bracts, without peduncular bracts), glabrous ovaries (vs. pubescent), and smaller, smooth-walled capsules (vs. mostly 6-9 mm long, with warty-roughened walls). *Sisyrinchium microbracteatum* is restricted to gypsum outcrops in central and south-central Nuevo León; *S. tenuifolium* apparently is much less common but has been collected in the same area.

Over almost all of its range, *Sisyrinchium tenuifolium* produces pubescent ovaries that mature into capsules with the surfaces conspicuously roughened by persistent, expanded trichome bases. Some plants of *S. tenuifolium* in Nuevo León and Coahuila are typical of the species in this respect, but in both states, other plants have glabrous ovaries and smooth fruits. In a population sampled from central Coahuila (*Wendt 1842* [ASU,TEX]), some plants have glabrous ovaries, others pubescent, but other differences among these plants are not apparent. Other accepted Mexican species very closely related to *S. tenuifolium*, however, are consistently separable from it with the emphasis of glabrous ovaries (e.g., *S. tolucense* Peyr., *S. palmeri* Greenm., *S. conzattii* Cald. & Rzed. - see Calderón 1988; McVaugh 1989), and when more specimens are at hand, the "*S. tenuifolium* complex" in northern México should be reexamined.

With the addition of the two new species described here, a total of six species of *Sisyrinchium* subg. *Echthronema* are presently known from the northeastern Mexican states of Coahuila, Nuevo León, and Tamaulipas:

Sisyrinchium microbracteatum and *S. hintoniorum* (Nuevo León),

Sisyrinchium arizonicum Rothr. (rare in Coahuila and Nuevo León),

Sisyrinchium convolutum Nocca (rare in Coahuila and Nuevo León),

Sisyrinchium tenuifolium Willd. (scattered in Coahuila and Nuevo León), and

Sisyrinchium schaffneri S. Wats. (common in Coahuila and Nuevo León, rare in Tamaulipas).

ACKNOWLEDGMENTS

I thank Billie Turner and Mark Mayfield for their review of the manuscript.

LITERATURE CITED

- Calderón de Rzedowski, G. 1988. Algunas consideraciones acerca de la familia Iridaceae en El Valle de México. *Acta Bot. Mex.* 1:21-27.
- McVaugh, R. 1989. *Sisyrinchium*. *Flora Novo-Galiciana* 15:311-330. University of Michigan Press, Ann Arbor, Michigan.
- Nesom, G.L. & L. Hernández S. 1992. A new species of *Sisyrinchium* subg. *Sisyrinchium* (Iridaceae) from México. *Phytologia* 73:429-434.

ADDITIONS TO THE NATIVE AND ADVENTIVE FLORA OF NEW MEXICO

Robert Sivinski

New Mexico Forestry and Resources Conservation Division, P.O. Box 1948,
Santa Fe, New Mexico 87504 U.S.A.

Timothy Lowrey

UNM Herbarium, Department of Biology, University of New Mexico,
Albuquerque, New Mexico 87131 U.S.A.

Roger Peterson

New Mexico Natural History Institute, P.O. Box 9314, Santa Fe, New Mexico
87504 U.S.A.

ABSTRACT

Sixteen new records of angiosperms are added to the New Mexico flora. Eight are native species and eight are adventive.

KEY WORDS: Flora, New Mexico

Botanical field surveys and herbarium searches by the authors have resulted in the following sixteen new records for the native and adventive flora of New Mexico. Several new angiosperm records for New Mexico have been recently published for the southern part of the state (Spellenberg *et al.* 1993). The report presented here is a more general accumulation of New Mexico records and has been produced to assist in preparing distribution maps for the remaining volumes of the *Flora of North America*. Nomenclature follows references cited.

AMARANTHACEAE

Amaranthus crassipes Schlecht. Chaves Co.: Bitter Lake National Wildlife Refuge (Salt Creek Wilderness), T8S, R25E, sect. 25, occasional in depressions in *Sporobolus airoides* grassland, elev. 1100 m, 30 October 1983, *Peterson 83-486* (UNM). Otero Co.: Upper Dog Canyon, T24S, R20E, section 25, sprawling on sand among matted *Bouteloua barbata*, elev. 1600 m, 23 October 1984, *Peterson 84-642* (UNM).

Comment: A native of tropical America that has become naturalized at eastern seaports. It was previously noted as far west as Trans-Pecos Texas (Correll & Johnston 1970).

APOCYNACEAE

Amsonia jonesii Woods. San Juan Co.: 12.2 road-kilometers east of Newcomb, west of confluence of Hunter's Wash and Chaco River on Pictured Cliff Sandstone, T24N, R16E, elev. 1670 m, 26 August 1961 (fruiting), *Cannon s.n.* (UNM); same location, 9 May 1962 (flowering), *Cannon s.n.* (UNM).

Comment: These collections represent a minor southern and eastern range extension from the adjacent states of Arizona, Colorado, and Utah.

ASTERACEAE

Matricaria matricarioides (Less.) Porter. Sandoval Co.: Jemez Mts., Capulin Canyon, Forest Road 501 on gravel roadway and adjacent road reclamation, T18N, R5E, sect. 22, elev. 2270 m, 5 June 1991, *Peterson 91-168* (UNM).

Comment: Martin & Hutchins (1981) included this species in *A Flora of New Mexico*, but with no certain records. This collection documents its residence in the state. It was observed again at the same location in 1993 and was spreading away from the gravel road. The reclamation of adjacent road disturbance occurred in 1989.

Onopordum acanthium L. San Miguel Co.: Pecos National Historical Park, 2.5 km south of Pecos on Hwy 63, T15N, R12E, sections 4, 5, 8, 9. and 16, in piñon-juniper woodland, common and well established on disturbed areas, elev. 2080 m, 30 July 1993, *Sivinski 2476* (UNM); same location, 3 July 1994, *Sivinski 2766* (NMC).

Comment: This European thistle is rapidly spreading throughout the western states. It is well established in disturbed areas on the highway, farms, and ranching operations near Pecos.

Scorzonera laciniata L. Bernalillo Co.: Albuquerque, University of New Mexico Campus, weed in flower bed near Geology Building, 13 May 1994,

Lowrey 1555 (UC). McKinley Co.: Red Rock State Park, 9 km E of Gallup, T15N, R17W, sect. 11, abundant in campground on deep sandy soil with *Bromus tectorum* and *Hordeum*, elev. 2020 m, 25 May 1994, *Sivinski & Lightfoot 2694* (NMC,UNM); Cañon de las Lagunitas, T16N, R5W, in *Bouteloua gracilis-Hilaria jamesii* rangeland, elev. 1950-2000 m, 11 June 1994, *Peterson 94-84* (UNM). San Miguel Co.: Las Vegas, weed in sidewalk, 29 May 1991, *Worthington 20939* (NMC); Pecos National Historical Park, disturbed area behind Trading Post stables at Fork Lightning Ranch, T15N, R12E, sect. 9 SW $\frac{1}{4}$, old landfill in piñon-juniper woodland, elev. 2070 m, 14 June 1993, *Sivinski & Simpson 2251* (UNM). Santa Fe Co.: near Edgewood, grassland just below level of juniper, May 1987, *Townsend s.n.* (NMC) [determined by Guy Nesom 1987]; Santa Fe, T17N, R10E, sect. 31, with *Bouteloua gracilis* and *Stipa comata* in *Pinus edulis* woodland, elev. 2150 m, 17 May 1993, *Peterson 93-139* (UNM).

Comment: This European species was first reported in North America along the eastern slope of Colorado in the 1950s (Great Plains Flora Assoc. 1986). It has since spread to Kansas and is now a common weed in north and west-central New Mexico. Leaf shape is quite variable and several of the New Mexican collections have narrower leaves than the description in the *Flora of the Great Plains* (Great Plains Flora Assoc. 1986). However, *Flora Europaea* (Tutin et al. 1976) describes this species as extremely variable in leaf shape.

BORAGINACEAE

Antiphytum floribundum (Torr.) A. Gray. Hidalgo Co.: Animas Mts., T32S, R19W, sect. 8 NW $\frac{1}{4}$, scattered and uncommon on a steep, S-facing slope of rocky rubble at the base of a rhyolitic cliff with *Quercus arizonica*, *Gymnosperma glutinosum*, and *Dasyilirion wheeleri*, elev. 2050 m, 21 August 1993, *Sivinski & Irick 2526* (BRY,MO,NMC,UNM).

Comment: First New Mexico record for the genus and species. This species ranges from the Trans-Pecos region of Texas south through the Sierra Madre Occidentale to central México.

Heliotropium fruticosum L. Grant Co.: City of Rocks State Park, T20S, R11W, sect. 4 SE $\frac{1}{4}$, abundant in shallow pockets of soil on large outcrops of rhyolitic rock, desert grassland of *Bouteloua eriopoda* and *Yucca elata*, with *Quercus emoryi* and *Juniperus deppeana* growing against the rock faces, elev. 1600 m, 16 October 1993, *Sivinski 2616* (BRY,NMC,UNM).

Comment: This annual ranges south to northern South America and east to the West Indies. It is a highly variable species with forms that are short and basally branched with small corollas, or tall and terminally branched with large corollas. The New Mexico collection belongs to the autogamous variant with minute (0.5-1 mm), included corollas and several short, basal branches, which is similar to the form in adjacent, southern Arizona.

BRASSICACEAE

Alyssum desertorum Stapf. McKinley Co.: Red Rock State Park just east of Gallup, growing in campground with *Ranunculus testiculatus*, 5 May 1980, *McCallum 699* (UNM); same location on deep, sandy soil, elev. 2040 m, 25 May 1994, *Sivinski & Lightfoot 2695* (NMC,UNM).

Comment: This European species is persistent, but not yet common in New Mexico. The first specimen was identified as *Alyssum alyssoides* (L.) L. Nevertheless, its glabrous silicles and early deciduous sepals clearly place it within *A. desertorum*. This species was noted to be not abundant at the time of the second collection.

Cardaria chalepensis (L.) Handel-Mazzetti. Bernalillo Co.: Albuquerque, in valley along roadside, 6 May 1951, *Clark s.n.* (UNM). Sandoval Co.: roadside close to Bernalillo, elev. 1550 m, 8 May 1931, *Castetter 4569* (UNM) [annotated by Ronald McGregor 1984].

Comment: This European species has not been recollected in New Mexico for more than forty years. It did occur in central Rio Grande Valley for at least twenty years and may still persist in that location.

EUPHORBIACEAE

Euphorbia carunculata Waterfall. Chaves Co.: Mescalero Sands, North Dunes 56 km east of Roswell, T11S, R31E sect. 6, elev. 1250 m, 13 September 1987, *Peterson 87-97* (UNM); Mescalero Sands, North Dunes 56 km east of Roswell and 1.6 km south of US 380, T10S, R30E, sect. 35, elev. 1240 m, prostrate on sand dunes partly anchored by *Quercus havardii*, 29 September 1991, *Peterson 91-401* (UNM).

Comment: This annual ranges from Oklahoma south to Chihuahua. It is locally common on sand dunes in this region (Great Plains Flora Assoc. 1986).

LAMIACEAE

Salvia pratensis L. Sandoval Co.: Near entrance to Jemez Falls Campground on Hwy 4, 2 October 1980, *Threlkeld s.n.* (UNM). Los Alamos/ Sandoval Co.: State Rt. 502 at county line, roadside, several clumps, 10 June 1990, *Foxx s.n.* (UNM).

Comment: These are the first records in the southwestern United States for this European species. It is an established weed in the Pacific Northwest east of the Cascade crests (Hitchcock & Cronquist 1973). The Fenton Lake, New Mexico collection label documents approximately 50 rosettes persisting for the previous three years. It was found a decade later as a roadside weed

further to the west of the first location. It is, therefore, established in the Jemez Mountains where it appears to be spreading.

MORACEAE

Morus alba L. Bernalillo Co.: Albuquerque, Rio Grande valley at Central Ave. Bridge, common in riparian woodland with *Populus fremontii* and *Ulmus pumila*, elev. 1510 m, 25 May 1994, Sivinski 2696 (NMC,UNM).

Comment: This Asian tree has been cultivated in the Rio Grande valley since at least 1906 (*Wootton s.n.* [NMC]). It is a well known feral tree, but this is apparently the first published record of its naturalization to New Mexico. The native cottonwood riparian forest of the Rio Grande valley through Albuquerque is frequently burned by wildfire. The fire replacement stand consists of *Elaeagnus angustifolia* L., *Ulmus pumila* L., *Morus alba*, and *Tamarix ramosissima* L. This stand of exotic trees is slowly replacing the native cottonwood bosque for several miles along the central portion of the Rio Grande.

POLEMONIACEAE

Gilia stellata Heller. Grant Co.: Gila National Forest, south-facing slopes above Middle Box of the Gila River, T18S, R18W, sect. 13, occasional in desert scrub dominated by *Mimosa biuncifera*, elev. 1360 m, 16 March 1983, Peterson 89-9 (UNM).

Comments: This species was previously known from the Mohave and Colorado Deserts. The southwestern New Mexico collection represents a significant eastward range extension from central Arizona.

RUBIACEAE

Galium bifolium S. Watson. San Juan Co.: Lukachukai Mts. (Chuska Mts.), 1 June 1950, Clark 15313 (UNM).

Comment: This species ranges from British Columbia to northwestern New Mexico and was included in the *Flora of New Mexico* (Martin & Hutchins 1981) as a single occurrence in Sandoval County. The *Flora* record, however, was based upon a misidentified specimen. The San Juan County collection correctly places this species in New Mexico.

Kelloggia galioides Torr. San Juan Co.: Lukachukai Mts., on old logging spur Navajo Forestry Road 7051I, 36°23'15" 109°02'45", ponderosa pine zone, white, sandy soil, elev. 2570 m, 22 June 1989, Hevron 184 (UNM).

Comments: First New Mexico record for the genus and species, which ranges from Idaho to northwestern New Mexico. The Lukachukai Mts. straddle the NM/AZ border and are often called the Chuska Mts. The presence of

this species and *Galium bifolium* represent an element of a more northwestern flora in this mountain range.

VIOLACEAE

Viola nuttallii Pursh. Colfax Co.: Above Sugarite Canyon in what is now Sugarite Canyon State Park, west of Lake Maloya near Segerstrom Creek, T32N, R24E, with grasses and forbs near *Pinus ponderosa* and *Quercus gambelii*, elev. 2340 m, 11 June 1984, *Peterson 84-299* (UNM). San Juan Co.: Mesa east of Los Pinos arm of Navajo Lake, T32N R7W, sect. 17, on sandy clay soil with *Artemisia tridentata* and *Pinus edulis*, elev. 1950 m, 9 May 1991, *Sivinski 1659* (NMC, UNM).

Comment: Martin & Hutchins (1981) included this species in *A Flora of New Mexico*, but with no certain records. These collections document the species for the state.

ACKNOWLEDGMENTS

This study was partially supported by a grant from the National Park Service, Pecos National Historical Park. Publication costs were paid by the New Mexico Energy, Minerals and Natural Resources Department, Forestry and Resources Conservation Division. The authors appreciate the manuscript review and comments of Richard Spellenberg and Kelly Allred. We also thank Helen Cannon and Bill Hevron for permission to publish their collection records, and John Strother (UC) for confirming our identifications of *Scorzonera laciniata*.

LITERATURE CITED

- Correll, D.S. & M.C. Johnston. 1970. *Manual of the Vascular Plants of Texas*. Texas Research Foundation, Renner, Texas.
- Great Plains Flora Assoc. 1986. *Flora of the Great Plains*. University Press of Kansas. Lawrence, Kansas
- Hitchcock, C.L. & A. Cronquist. 1973. *Flora of the Pacific Northwest*. University of Washington Press. Seattle, Washington.
- Martin, W.C. & C.R. Hutchins. 1980, 1981 (vols. 1 & 2 respectively). *A Flora of New Mexico*. J. Cramer, Vaduz, Liechtenstein.

- Spellenberg, R., L. McIntosh, L. Brouillet. 1993. New records of angiosperms from southern New Mexico. *Phytologia* 75:224-230.
- Tutin, T.G., V.H. Heywood, N.A. Burges, D.M. Moore, D.H. Valentine, S.M. Walters, D.A. Webb. 1976. *Flora Europaea*. Vol. 4, Cambridge University Press, Cambridge, Great Britain.

BOOKS RECEIVED

Saltmarsh Ecology Paul Adam. Cambridge Studies in Ecology. Cambridge University Press, 40 West 20th Street, New York, New York 10011-4211. 1993. xii. 461 pp. \$37.95 ISBN 0-521-44823-9 (paper).

This paperback edition of a volume first published in 1990 provides a broadly comprehensive view of saltmarshes. Although much of the text is drawn from saltmarshes of the British Isles, considerable information is also drawn from Australia, with somewhat less coverage of saltmarshes in other parts of the world. As might be expected most of the contents deal with coastal salt marshes, but inland saltmarshes are discussed also. The introductory chapter examines general features of saltmarshes, including processes contributing to their formation, and defines what constitutes a saltmarsh for the purpose of this book. Subsequent chapters examine the biota, types and causes of variation, environmental factors affecting saltmarshes, human effects on saltmarshes, and other topics.

The Cruciferae of Continental North America, Systematics of the Mustard Family from the Arctic to Panama. Reed C. Rollins. Stanford University Press, Stanford, California 94305-2235. 1993. xviii. 976 pp. \$125.00 ISBN 0-8047-2064-9 (cloth).

This massive compendium on the mustards of North America is an essential component of the library of any researcher planning a study of members of the family. The bulk of the content consists of keys to and descriptions of genera, species, and varieties. Black and white photographs and line drawings supplement the text. The only negative comment might be that the book includes several new combinations and taxonomic descriptions. These might best have been first published elsewhere, then included in the book.

The Masked Bobwhite Rides Again. John Alcock. The University of Arizona Press, 1230 N. Park Avenue, Suite 102, Tucson, Arizona 85719-4140. 1993. xii. 186 pp. \$16.95 ISBN 0-8165-1405-4 (paper); \$35.00 ISBN 0-8165-1387-2 (cloth).

This book is the third by the same author in which he provides ecological insight to the general public through a series of essays describing natural systems. In the case of the present volume, the main topic is the deserts of the southwestern United States, but Alcock uses everything from ants and Apaches, to coyotes and towhees to bring interesting features of the desert to the lives of readers who may never have seen a desert. The book is interesting and thought provoking not only for the general public, but for scientists as well.

The Message of Fossils. Pascal Tassy. McGraw-Hill, Inc., 11 West 19th Street, New York, New York 10011. 1993. 163 pp. \$10.95 ISBN 0-07-062947-1 (paper).

Published through McGraw-Hill's *Horizons of Science Series*, this book examines the concept of evolution more than fossilization *per se*. The impacts of fossils on Darwin's theory, as well as modifications of his ideas and more modern ideas of evolutionary mechanisms are discussed.

INDEX TO AUTHORS, VOLUME 76

- Babu, G.J. 461
Cuatrecasas, J. 402
Gandhi, K.N. 441
Glad, J.B. 307
Grant, J.R. 284
Grant, V. 303
Halse, R.R. 307
Hilu, K.W. 410
Kartesz, J.T. 441
King, R.M. 14
Landry, P. 73
Lowrey, T. 473
McIntosh, L. 329
McMullen, C.K. 30
Mendenhall, M.G. 383
Mishaga, R. 307
Naidu, K.C. 461
Nesom, G.L. 85, 96, 101, 106, 115,
125, 136, 143, 160, 168, 176,
185, 193, 406, 426, 465, 468
Palacios C., R. 433
Peterson, R. 473
Powell, A.M. 324
Pushpalatha, V. 461
Quiroz G., D.L. 433
Reveal, J.L. 1
Robinson, H. 14, 19, 24, 27
Sivinski, R. 473
Turner, B.L. 8, 39, 52, 69, 277, 281,
290, 333, 345, 385, 389, 391,
412, 421
Viderman, D.M. 30
Yarborough, S.C. 324
Zanowiak, D. 106

INDEX TO TAXA, VOLUME 76

New taxa described in this volume are indicated in bold face type.

- Acalypha* 463
 indica 463
Acamptopappus 211, 271
Acanthaceae 433, 439, 440, 461
Acantheae 439
Acanthospermum 462
 hispidum 462
Achaetogeron 272
Achnophora 197, 207, 229, 230, 273
Achyranthes 462
 aspera 462
Acorus 2, 6
 calamus 6
Acrocephalus 463
 indicus 463
Adenostemma 14-16, 390
 flintii 16
 harlingii 14
 platyphyllum 15
 zakii 14, 15
Aerva 462
 lanata 462
Agathaea 93
 corymbosum 93
Agave 66
 lechugilla 66
Ageratum 462
 conyzoides 462
Agrostis 312
Aizoaceae 462
Alismatidae 2
Allmania 462
 nodiflora 462
Allophyllum 305
Almutaster 212, 271
Alternanthera 462
 sessilis 462
Alysicarpus 463
 monilifer 463
Alyssum 476
 alyssoides 476
 desertorum 476
Amaranthaceae 462, 474
Amaranthus 462, 474
 crassipes 474
 spinusus 462
Amborellaceae 3
Ambrosia 329, 330
 psilostachya 329, 330
Amellus 85, 87, 88, 90, 195, 197,
 200, 205, 218, 219, 223-225,
 251, 269, 274
Amphiachyris 211, 271
Amphipappus 211, 237, 271
Amsonia 474
 jonesii 474
Anaphalis 187, 188, 191
Anazeton 187
Andrographis 461
 echioides 461
 paniculata 461
Angianthinae 242
Anisochilus 463
 carnosus 463
Anisomeles 463
 indica 463
Antennaria 187, 188, 403
 andina 403
Anthemideae 196, 226, 241, 248, 260,
 263

- Antiphytum* 475
 floribundum 475
Aphanostephus 203, 216, 218, 251,
 262, 270, 271
 ramosissimus 216
 skirrhobasis 262
 Apiaceae 442
Apluda 463
 mutica 463
 Apocynaceae 462, 474
Apodocephala 90, 243
Apopyros 122, 124, 168-170, 174,
 176, 177-183, 204, 220, 267,
 272
 corymbosus 176, 178-180, 182
 warmingii 176, 177, 179-181,
 183
Apostates 163, 166, 242, 243, 264
 Araliales 5
 Aralianae 5
 Aralidiales 5
Arbutus 250, 365
Archibaccharis 109, 199, 202, 247,
 271
Arctogeron 201, 241, 273
 Arecidae 2
Argemone 463
 mexicana 463
Aristida 463
 funiculata 463
 Aristolochiaceae 3
 Aristolochiales 3
Artemisia 478
 tridentata 478
Arnica 91
 crenata 91
Asanthus 389, 390
 Asclepiadaceae 462
Asiris 272
Aspilia 24, 25
 retroflexa 24, 25
Aster 90-92, 95, 105, 115, 117, 120,
 121, 173, 174, 176, 178, 179,
 182, 193, 195, 201,
 212, 224, 240, 245, 251, 266-
 268, 273
 alpinus 245
 bakeranus 95
 camporum 115, 120, 121
 crenatus 91
 grisebachii 105, 174, 267
 harveyanus 90
 kingii 266
 longipes 117
 nanus 92
 Oxytripolium group 120
 subg. *Oxytripolium* 182
 perezoides 92
 subg. *Pteropappus* 91
 regnellii 182
 setosus 176, 178, 179, 182
 squamatus 182
 tazifolius 92
 tuberosus 176, 178, 179
 vahlkii 120, 121
 warmingii 176, 177
 Asteraceae 14, 18, 19, 23, 24, 26,
 27, 29-31, 39, 51, 85, 94-96,
 100, 101, 105, 106, 113-115,
 123-125, 134-136, 142, 143,
 158-160, 163, 166-168, 174-
 176, 184, 185, 191, 193, 245,
 246, 259-271, 281, 283, 324,
 389, 390, 405, 426, 457, 462,
 474
 Astereae 85, 94-96, 100, 101, 102,
 105, 106, 113-115, 122-126,
 131, 134-136, 142, 143, 147,
 152, 158-160, 163, 166-168,
 175, 176, 183, 184, 193-201,
 212, 213, 216, 218, 219, 222,
 223, 225, 226, 229, 231, 232,
 235, 236, 238, 239, 241-246,
 248, 250, 252, 260-271, 403,
 405, 426
 Asteridae 3, 5

- Asterinae* 193, 194, 199, 201, 212,
 215-217, 220, 240, 241, 243,
 245-247, 249, 251, 271-273
Asteroideae 94, 105, 113, 123, 183,
 188, 191, 196, 226, 229, 259,
 264
Asteromoea 273
Asteropsis 101, 103, 104, 106, 112,
 114, 115, 122, 124, 126, 172,
 175, 197, 210, 234, 235, 267,
 272
 macrocephala 103, 104, 112, 172,
 235
Asterothamnus 201, 273
Astranthium 203, 217, 251, 261, 271
Atylosia 463
 rugosa 463
 scarabaeoides 463
Avicenniaceae 344
Ayapana 14, 16-18, 390
 herrerae 14, 16, 17
Aylacophora 228, 273
Aztecaster 95, 135, 167, 207, 266,
 271

Baccharidastrum 112, 273
 notobellidiastrum 112
 rivulare 112
Baccharidinae 95, 109, 114, 135, 167,
 194, 197-199, 202, 213, 214,
 227, 228, 239, 245-249, 254,
 259, 266, 271, 273, 274
Baccharidopsis 273
Baccharis 109, 193, 198, 202, 213,
 214, 220, 227, 247, 263, 273,
 403
 aretioides 403
 helichrysoides 214
Bacopa 463
 monnieri 463
Baptisia 383, 384
 australis 383
 var. **aberrans** 383

Baptisia (cont.)
 minor 383
 var. *aberrans* 383
Barleria 461
 strigosa 461
Bataceae 4
Batales 4
Bauhinia 333-343
 bartlettii 333, 334-337, 340, 342
 bifoliola 336
 congesta 337
 coulteri 333, 334, 336, 338
 var. *arborescens* 336
 var. *coulteri* 336
 divaricata 333, 334, 336, 339
 jermiana 337
 lunarioides 333, 334, 337, 341
 macranthera 333-337, 340, 342
 var. *grayana* 337
 monantha 333, 343
 purpusii 342
 ramosissima 333-335, 337, 341-
 343
 retifolia 337
 runyonii 342
 unguicularis 342
 uniflora 333, 334, 338, 342, 343
Begoniaceae 4
Bellidiastrum 215, 273
Bellidinae 193, 202, 215, 217, 248,
 273
Bellis 193, 194, 198, 200, 202, 212,
 215, 217, 224, 273
 perennis 215
 scapiformis 217
Bellium 193, 205, 215, 224, 273
Belloa 187
Benitoa 209, 233, 234, 264, 271
Bigelowia 211, 271
Bignoniaceae 259
Biota 272
Bisgoeppertia 8
Blakiella 101, 103, 106, 112, 115,

- Blakiella* (cont.) 122, 126, 197, 210,
 227, 234, 235, 273
 bartsiiifolia 112
Blastania 441, 442, 444
 cerasiformis 444
 fimbristipula 444
Blepharis 461
 asperrima 461
Boltonia 193, 201, 212, 216, 218,
 225, 240, 245, 271
 Boraginaceae 462, 475
Borreria 463
 articularis 463
 stricta 463
Bouteloua 474, 475
 barbata 474
 eriopoda 475
 gracilis 475
Brachiaria 463
 distachya 463
Brachyactis 201, 251, 272, 273
Brachychaeta 272
Brachycome 128, 136-138, 141, 142,
 198, 200, 203, 216, 217, 219,
 231, 261, 273
 cardiocarpa 141
 neocaledonica 136, 138, 141
 nivalis 141, 217
 readeri 216
 sarasinii 136, 138
 scapiformis 141
 scapigera 141
Brachycominae 193, 198, 200, **203**,
 215-219, 223, 232, 248, 250,
 251, 257, 271-273
Bradburia 266, 272
 Brassicaceae 4, 442, 443, 476
 Brexiales 4
Brickellia 390
 fendleri 390
Brickelliastrum 14, 17, 18, 389, 390
 fendleri 18, 389, 390
 villarealii 14, **17**, 389, 390
Brintonia 272
 Bromeliaceae 284, 289
Bromus 475
 tectorum 475
Bryonopsis 462
 laciniosa 462
Buettneria 464
 herbacea 464
 Byblidales 5
Cacalia 404
 ilicifolia 404
 Caesalpinaceae 462
 Caesalpinieae 333
 Caesalpinioideae 343
Caesulia 462
 azillaris 462
Callistephus 197, 201, 273
Calotis 141, 197, 198, 200, 203, 216,
 217-219, 227, 231, 261, 273
 anamitica 217, 218
 caespitosa 218
 gaudichaudii 218
Calotropis 462
 gigantea 462
Caltha 441, 442, 445, 446, 456
 flabellifolia 445, 446
 palustris 441, 445, 446
 var. *flabellifolia* 441, 445, 446
 var. *palustris* 445, 446
 var. *radicans* 445, 446
 radicans 446
Calypocarpus 40, 50
 vialis 50
 wendlandii 50
Camptacra 125, 126, 128-131, 133,
 134, 144, 146, 147, 157, 210,
 260, 273
 Capnanthemae 3
 Capparaceae 4
 Capparales 3, 4
Capparanae 1, **3**
 Capparidaceae 462

- Capparis* 462
 zeylanica 462
 Caprifoliales 5
Cardaria 476
 chalepensis 476
Cardiospermum 463
 halicacabum 463
Carex 321
 Caricaceae 3, 4
 Caricales 3, 4
 Caryophyllidae 3
Casparia 337, 342, 343
 congesta 337
 jermiyana 337
 lunarioides 337
 macranthera 337
 monantha 343
 purpusii 342
 ramosissima 342
 retifolia 337
 runyonii 342
 unguicularis 342
 uniflora 343
Cassia 462
 absus 462
 occidentalis 462
 sophora 462
 tora 462
Cassinia 188
 Cassiniinae 187, 188
Castilleja 329-332, 406-409, 465-467
 aspera 465, 466
 sect. *Castilleja* 408, 409
 subg. *Castilleja* 408, 409
 chlorosceptron 465, 466
 durangensis 465, 466
 sect. *Epichroma* 408, 409
 sect. *Euchroma* 330, 466
 fissifolia 408
 glandulosa 329, 330
 gonzalezii 465
 lebgueana 406-409
 macrostigma 329, 330
Castilleja (cont.)
 ornata 329-331
 ortegae 406-409
 sect. **Ortegae** 406, 409
 Ortegae group 406-408
 palmeri 329, 330
 pediaca 329, 330
 pteroaulon 407, 409
 scorzoneraefolia 330
 sphaerostigma 329, 330
 tayloriorum 408
 tenuiflora 409
 Tenuiflorae group 408
 Castillejinae 409
Catharanthus 462
 pusillus 462
 roseus 462
Cavea 265
 Celastranae 4
Celmisia 155, 159, 160, 165, 166,
 207, 228, 229, 262, 273
 Censitae 420, 425
Centaurium 8
Centipeda 200, 206, 225-227, 248,
 273
 cunninghamii 226
 minuta 226
 brachycomoides 157
 sect. *Lignosae* 165
Ceratogyne 141, 203, 216, 217, 273
Ceratophyllum 2
 Cercideae 343
Cercis 370
Ceruana 197, 206, 274
Chaenomeles 441, 442, 451, 457
Chaetopappa 197, 199, 205, 206, 223-
 225, 241, 250, 266, 269-271
 bellioides 224
 elegans 266
Chamaegeron 205, 223, 273
Charieis 88
 heterophylla 88
Chevreulia 187

- Chiliophyllum* 207, 251, 273
Chiliotrichopsis 207, 251, 273
Chiliotrichum 195, 207, 251, 273
Chionolaena 185-191
 concinna 190
 eleagnoides 189
 mexicana 185, 189, 190
 seemannii 185, 186
Chlamyditis 273
Chloracantha 212, 271
 Chloranthaceae 3, 6
 Chloranthales 3
Choenomeles 441, 442, 451
Chrozophora 463
 rotleri 463
Chrysanthemum 241
Chrysocoma 87, 198, 205, 274
Chrysoma 211, 271
Chrysopsidinae 193, 198, 199, **203**,
 219, 220, 233, 240, 241, 245,
 249, 272
Chrysopsis 169, 173, 195, 197, 203,
 204, 266, 269, 272
Chrysothamnus 211, 237, 238, 268,
 272, 273
Cicenda 8
 Cirrhatae 4
Cirsium 312
Cleome 462
 aspera 462
 monophylla 462
 viscosa 462
Colobanthera 206, 225, 274
Columbiadoria 237, 266, 272
 Commelinidae 2
Commidendron 202, 213, 214, 247,
 274
 robustum 214
Complaya 26
 Compositae 18, 26, 50, 94, 100, 105,
 113, 114, 123, 124, 134, 142,
 158, 159, 166, 167, 174, 175,
 183, 191, 259-266, 268-271,
 Compositae (cont.) 328, 402, 405
Coincya 441-443, 456
 monensis 441, 443
 ssp. *recurvata* 443
 var. *recurvata* 441, 443
Coldenia 462
 procumbens 462
Commelina 462
 attenuata 462
 bengalensis 462
 erecta 462
 longifolia 462
 Commelinaceae 462
 Convolvulaceae 457, 462
Conyza 104-109, 112, 122, 124, 128,
 143, 157, 158, 170, 174, 176,
 178, 179, 182, 193, 195, 200,
 204, 206, 213, 220, 222, 228,
 234, 235, 238, 248, 250, 251,
 258, 266, 268, 271, 273
 blanchetii 176, 178, 179, 182
 bonariensis 109, 157
 canadensis 108, 157, 222
 notobellidiastrum 106, 107, 109,
 112
 primulifolia 108
 rivularis 106, 107, 109, 112
 setosa 178, 179
 spinulosa 104
 trihecatactis 108
Conyzanthus 272
 Conyzinae 193, 199, 204, 220, 235,
 245, 246, 248, 250, 251, 258,
 272, 273
Corchorus 464
 aestuans 464
 Coreopsidinae 163, 269
Corethrogyne 195, 197, 209, 233, 234,
 272
 Coriariaceae 4
 Coriariales 4
 Cornaceae 457
 Cornales 5

- Cornanae 4, 5
 Corniculatae 4
Cornidae 1, 3, 4, 5
Cornus 7, 450, 454
 citrifolia 450, 454
 Cotuleae 260
 Crassulaceae 4
Crinitaria 201, 212, 251, 273
Critoniopsis 27
 aristeguietae 27
Croptilon 204, 233, 272
 Crossosomataceae 4
Crotalaria 463
 linifolia 463
 nana 463
 prostrata 463
 verrucosa 463
Croton 463
 bonplandianum 463
 Cruciferae 456, 479
Ctenolepis 441, 442, 444, 462
 cerasiformis 444
 garcini 462
Cuatrecasanthus 29
Cuatrecasasiella 187
Cucumis 462
 trigonus 462
 Cucurbitaceae 4, 442, 444, 456, 462
 Cucurbitales 4
Cucurbitanae 1, 4
Culcitium 404
 pavonii 404
Cyanotis 462
 azillaris 462
Cyathocline 196, 206, 225, 227, 248,
 273
Cynodon 463
 dactylon 463
 Cyperaceae 462
Cyperus 462
 aloppecuroides 462
 cephalotus 462
 compactus 462
 Cyperus (cont.)
 difformis 462
 distans 462
 iria 462
 platystylis 462
 triceps 462
Cytisus 444
 laburnum 444
Dactyloctenium 463
 aegyptium 463
Dalea 510
Damnamenia 159, 165, 166
Dacryotrichia 206, 225, 274
Damnamenia 207, 262, 273
Darwiniothamnus 30-36, 221, 222,
 272
 tenuifolius 30-36
Dasyilirion 475
 wheeleri 475
 Datisceae 4
Desmazieria 460
Desmodium 463
 gangetium 463
 triflorum 463
Dichaetophora 203, 272
Dichrocephala 206, 225, 226, 274
Dichromochlamys 126, 130, 132, 134,
 144, 146, 147, 210, 216, 234,
 236, 261, 273
 Dicrastylidaceae 344
Digera 462
 muricata 462
 Dillenidae 2, 3
Dimorphocoma 126, 130, 133, 144,
 147, 197, 210, 230, 237, 242,
 273
Diplopappus 171-173, 176, 178
 corymbosus 176, 178
 graminifolius 171
 pinifolius 172
 stenophyllus 171
 villosus 173

- Diplostegium* 166, 207, 214, 227,
 247, 251, 273
crassifolium 166
- Dipsacales 5
- Dipterocanthus* 461
prostratus 461
- Dodonaea* 463
viscosa 463
- Doellingeria* 212, 240, 267, 272
- Ducampopinus* 78
- Dyscritogyne* 389, 390
- Eastwoodia* 197, 211, 272
- Eclipta* 462
alba 462
- Ecliptinae 24, 26, 40
- Egletes* 200, 206, 216, 225, 226, 269,
 273
- Elachanthus* 126, 130, 133, 144, 147,
 197, 210, 230, 237, 242, 274
- Elaeagnus* 477
angustifolia 477
- Elaeocarpaceae 4
- Elaeocarpales 4
- Elaphandra* 24-26
retroflexa 25
- Eleocharis* 321
- Eleusine* 410, 411
africana 410, 411
coracana 410, 411
 subsp. **africana** 410, 411
indica 410, 411
 subsp. *africana* 410, 411
- Elphegea* 91, 92
ciliata 92
microcephala 92
 subg. *Zyrphelis* 91
- Embryophyta 142
- Emilia* 462
sonchifolia 462
- Engleria* 87, 205, 222, 274
- Eragrostiella* 463
bifaria 463
- Eragrostis* 463
tenella 463
unioloides 463
- Eremiastrum* 272
- Ericameria* 193, 207, 221, 228, 246,
 248, 250, 253, 268, 272
- Erigeron* 96-100, 104, 109, 111, 112,
 117, 120, 122-123, 135, 142,
 143, 144, 146-148, 150-157,
 159, 160, 163, 165-167, 170-
 174, 178, 183, 184, 191, 193-
 195, 198, 200, 204, 213, 216,
 220-222, 235, 238, 246, 248,
 250, 251, 263-268, 270, 272,
 426
- acris* 222
- ambiguus* 143, 144, 146, 147
- blanchetii* 178
- bonariensis* 157
- brachycomoides* 157
- calcicola* 99
- camporum* 120
- canadensis* 157, 222
- candollei* 157
- chiangii* 96, 99, 100
 var. *chiangii* 99
- sect. *Cincinnati* 222
- conyzoides* 143, 156, 157
- coronarius* 266
- crispus* 157
- decurrens* 157
- dianthifolius* 171
- divergens* 99
- dubius* 172
- sect. *Erigeron* 221
- filiformis* 171
- flagellaris* 99
- gardneri* 112
- sect. *Geniculactis* 266
- glaucifolius* 173
- gunnii* 154
 var. *bellidioides* 154
- hieracifolius* 111

Erigeron (cont.)

- sect. *Karvinskia* 99
- karvinskianus* 143
- sect. **Lamprocaules** 96, 98, **99**
- liatroides* 157
- sect. *Linearifolii* 99, 100, 266
- minurioides* 157
- montevidensis* 171, 173
- mucronatus* 143
- nudiscapus* 117
- modestus* 97
- sect. *Olygotrichium* 97, 216
- pappochroma* 426
- pappocromus* 150-156, 165
 - var. *billardierei* 154
 - var. *gunnii* 154
 - var. *oblongatus* 154
 - var. *pappocromus* 154
 - var. *setosus* 150
 - var. *stellatus* 151
- pattersonii** 96, 97, 99, 268
- paucifolius* 112
- phlogotrichus* 154
- pinifolius* 171
- sect. *Polyactis* 266
- primulifolia* 111
- pubescens* 99
- rapensis* 160, 163
- resinosus* 172
- rivularis* 112
- scoparioides* 96, 99
- sessilifolius* 144, 146, 147
- setosus* 148, 150, 152
- stellatus* 148, 151, 152
- strigosus* 109
- tasmanicus* 151, 154
- var. *stellatus* 151
- tenuis* 109
- tridactylus* 104
- trifurcatus* 104
- trilobus* 157
- subg. *Trimorpha* 204, 221, 222, 238, 248

Erigeron (cont.)

- tweediei* 183
- uniflorus* 221
- vittadinia* 157
- Eriocarpum* 272
- Eriocaulaceae 344
- Erodiophyllum* 197, 206, 225, 227, 248, 274
- Erythanthera* 159
- Erythriinae 8
- Eucephalus* 212, 219, 272
- Eupatorieae 14, 18, 389, 390
- Eupatorium* 18, 389, 390
 - coelocaula* 18
 - fendleri* 389, 390
- Euphorbia* 325, 463, 476
 - carunculata* 476
 - hirta* 463
- Euphorbiaceae 463, 476
- Eurybiopsis* 125, 126, 134, 260, 274
 - macrorhiza* 125
- Euthamia* 211, 237, 250, 268, 272
- Ewartia* 187, 188
- Fabaceae 290, 343, 383, 385, 387, 442, 444, 457, 463, 510
- Fabanae 510
- Facelis* 187
- Felicia* 85, 87-90, 92-94, 157, 198, 199, 205, 223, 225, 262, 274
 - ciliaris* 93
 - sect. *Dracontium* 94
 - heterophylla* 88
 - lasiocarpa* 92
 - ovata* 94
 - tenella* 157
- Feliciinae** 193, 197-200, **205**, 215, 218, 219, 222-225, 244, 248, 250, 257, 271, 272
- Ferreyranthus* 19
 - gentryi** **19**
- Festuca* 312
- Fimbristylis* 462

- Fimbristylis* (cont.)
dichotoma 462
ovata 462
- Fleischmannia* 17
sideritidis 17
- Floscaldasia* 207, 225, 228, 267, 273
- Flosmutisia* 207, 228, 273
- Formania* 241, 269
mekongensis 241
- Francoaceae 4
- Frangula* 441, 442, 447-450, 453
betulifolia 441, 448
 ssp. *obovata* 441, 448
 × *blumeri* 441, 448
californica 441, 448, 449
 ssp. *crassifolia* 441, 448
 ssp. *cuspidata* 441, 449
 ssp. *occidentalis* 441, 449
 ssp. *tomentella* 441, 449
 ssp. *ursina* 441, 449
citrifolia 450
rubra 441, 449
 ssp. *modocensis* 441, 449
 ssp. *nevadensis* 441, 449
 ssp. *obtusissima* 441, 449
 ssp. *yosemitana* 441, 449
sphaerosperma 441, 450
 ssp. *longipes* 441, 450
- Frazinus* 309, 310, 321
latifolia 321
- Galatella* 197, 201, 212, 251, 273
- Galeobdolon* 454
- Galium* 463, 477, 478
asperifolium 463
bifolium 477, 478
- Garryales 5
- Geissolepis* 197, 203, 218, 269, 272
- Geniostemon* 8-13
atarjanum 8, 9, 10, 12
coulteri 8-10, 12
gypsophilum 8, 9, 11-13
schaffneri 8, 9, 11-13
- Gentianaceae 8, 13
- Gentiananae 5
- Gerbera* 91
crenata 91
- Gilia* 52-68, 303-306, 477
acerosa 66
 sect. *Arachnion* 52, 303, 304
austro-occidentalis 305
 subg. *Benthamiophila* 52
brecciarum 303-305
 subsp. *brecciarum* 303-305
 subsp. *jacens* 303, 305
campanulata 59
diegensis 304
filiformis 59
foetida 52, 59
 sect. *Gilia* 52
 subg. *Gilia* 52
 sect. *Giliandra* 52
 sect. *Giliastrum* 52, 53, 57, 59
 sect. *Gilmania* 52
glandulosa 66
gypsophila 52-54, 55, 56, 58, 67
incisa 52, 53, 56, 59, 60
insignis 52-54, 57, 59, 61
interior 305
jacens 303-305
latifolia 59
leptantha 52, 303
 subsp. *purpusii* 52
lindheimeriana 56
ludens 52-54, 56-58
palmeri 52, 54, 57, 61
 subsp. *spectabilis* 57
perennans 53, 56
platyloba 52, 53, 65, 67
purpusii 52-54, 57, 62, 65-67
rigidula 52-54, 56, 57, 59, 63, 65-67
 var. *acerosa* 52, 53, 56, 63, 65, 66
 subsp. *insignis* 52, 57

- Gilia rigidula* (cont.)
 subsp. *rigidula* 52
 var. *rigidula* 52, 57, 59, 63, 65-67
 sect. *Saltugilia* 52
 stellata 477
 stewartii 52-54, 64, 65, 67
 tenuiflora 303-305
 tenuiflora × *brecciarum* 303
- Giliastrum* 66
 acerosum 66
 rigidulum 66
- Giseckia* 462
 pharmacoides 462
- Glinus* 462
 lotoides 462
 oppositifolius 462
- Gnaphalieae 185, 191
- Gnaphaliinae 402
- Gnaphalinae 229, 242, 244, 248, 259
- Gnaphaliothamnus* 185-191
 concinus 185, 186, 189, 190
 costaricensis 186
 cryptocephalus 186
 durangensis 186
 eleagnoides 186, 189, 190
 lavandulifolius 186, 187
 salicifolius 185-187, 189, 190
 sartorii 186
- Gnaphalium* 189, 191, 229
 concinnum 189
 lavandulaceum 189
- Gomphrena* 462
 decumbens 462
- Goodeniaceae 246
- Gramineae 411
- Grangea* 128, 193, 195, 200, 206, 217, 225-227, 274
 maderaspatana 217
 zambesiaca 226
- Grangeinae 193, 194, 195, 198, 200, 206, 215-217, 219, 222, 225-227, 231, 232, 245-248, 255,
- Grangeinae (cont.) 262, 273, 274
- Grangeopsis* 200, 206, 274
- Grauanthus* 206, 274
- Greenella* 272
- Greyiaceae 4
- Grindelia* 195, 196, 209, 232, 233, 238, 247, 250, 260, 268, 272
- Grossulariaceae 4
- Grosvenoria* 14, 18, 390
 coelcaulis 18
 jelskii 18
- Gundlachia* 198, 211, 272
- Gutierrezia* 211, 237, 247, 250, 272
- Gymnaster* 273
- Gymnosperma* 211, 272, 475
 glutinatum 475
- Gymnostephium* 85-90, 92-94, 199, 205, 225, 274
 angustifolium 93
 ciliare 93
 corymbosum 85, 88, 93
 fruticosum 88, 93
 gracile 88, 93
 hirsutum 92, 93
 leve 88, 89, 93
 papposum 85, 93
- Gynandropsis* 462
 pentaphylla 462
- Gynozys* 404
 aquifolia 404
 ilicifolia 404
- Gyrodoma* 206, 225, 274
- Gyrostemonaceae 4
- Hamamelididae 2, 3
- Haplopappus* 111, 115-117, 151, 154, 171, 209, 232, 233, 240, 249, 250, 260, 263, 273, 426
 bellidioides 154
 blanchetianus 111
 gunnii 154, 426
 sect. *Gymnocoma* 260
 sect. *Inulopsis* 115, 116

- Haplopappus* (cont.)
marginatus 171
pappocromus 154
scaposa 115-117
stellatus 151
stoloniferus 116
tasmanicus 154, 426
- Hazardia* 209, 232, 260, 272
- Hedeoma* 277-280
jucundum 279
patrinum 277-279
 subg. *Poliomenthoides* 277
rzedowskii 277-279
- Heleastrum* 212, 240, 250, 272
 Helenieae 324, 328
- Heliantheae* 24, 26, 39, 40, 51, 163,
 191, 243, 264, 268
- Helichrysum* 188
- Helicteres* 464
isora 464
- Heliotropium* 462, 475
fruticosum 475
indicum 462
scabrum 462
- Heracleum* 441-443
lanatum 441-443
maximum 441-443
- Herrickia* 212, 272
- Hesperodoria* 238, 266, 272
- Heteractis* 93
falcata 93
- Heterochrominae 201, 211
- Heteromma* 206, 227, 274
- Heteropappus* 198, 201, 273
- Heteroplexis* 202, 213, 247, 260, 273
- Heterothalaminae 202
- Heterothalamus* 109, 202, 273
- Heterotheca* 204, 250, 269, 272
- Hilaria* 475
jamesii 475
- Hinterhubera* 206, 207, 235, 273
- Hinterhuberinae 85, 90, 95, 121, 130,
 135, 160, 165-167, 193, 194,
- Hinterhuberinae (cont.) 197-199, 206,
 214, 227-229, 241, 243, 245-
 251, 253, 266, 267, 271-274
- Holcus* 312
lanatus 312
- Homochroma* 85, 86, 89, 91, 211
angolensis 91
ecklonis 91
- Homochrominae 201, 210, 211, 273,
 274
- Hordeum* 475
- Hydrangeales 4, 5
- Hyptis* 463
suaveolens 463
- Hysterionica* 101, 103, 105, 115, 117,
 122, 124, 168-174, 176, 182,
 184, 198, 204, 220, 267, 273
- aberrans* 169, 173
 var. *aberrans* 173
 var. *hunzikeri* 173
- bakeri* 172, 173
 var. *aberrans* 173
- cabreriae* 173
- dianthifolia* 171, 172
 var. *pulvinata* 172
- filiiformis* 103, 171
- glaucifolia* 169, 173
- jasionoides* 173, 174
- linearifolia* 172
- marginata* 105, 171, 174, 267
- montevidensis* 169, 173
- pinifolia* 171
- pulchella* 173
- pulvinata* 172
- scaposa* 117
- setuligera* 171
- subvillosa* 172
- villosa* 173
- Illiciales 3
- Inula* 116
- Inuleae 185, 191, 226, 229, 238, 241,

- Inuleae* (cont.) 243, 248, 259, 262,
402, 403, 405
Inuloideae 199, 261
Inulopsis 101, 103, 113-122, 170, 174,
197, 210, 234, 235, 267, 273
 camporum 115-120, 121
 phoenix 115, 117, 118-121
 scaposa 115, 117-119
 stenophylla 115, 117-119
Iogeton 26
Ionactis 197, 212, 220, 240, 241, 249,
266, 268, 272
Iotasperma 126, 127, 130, 132, 143,
144-147, 210, 216, 237, 274
 australiensis 143-146
 sessilifolia 143, 146
Ipomoea 462
 carnea 462
 hederifolia 462
 nil 462
 pestigradis 462
Ipomopsis 68
Iridaceae 472, 468
Isocoma 209, 233, 272
Isoetopsis 126, 242, 269
 graminifolia 242
Ixioclamys 106, 113, 126, 130, 132,
134, 144, 146, 147, 197, 210,
234, 236, 261, 274
 cuneifolia 236
 filicifolia 236
 integerrima 236
 nana 236
Izodia 188

Jefea 26
Jeffreya 205, 274
Joseanthus 27, 29
 sparrei 27
 trichotomus 27
Juglans 370
Juncus 321, 330
 balticus 330

Juniperus 374, 475
 deppeana 475
Justicia 461
 glauca 461
 procumbens 461
 simplex 461

Kalimeris 200, 201, 216, 240, 273
Kelloggia 477
 galiioides 477
Kemulariella 201, 273
Keteleeria 78
Keysseria 141, 151, 156, 208, 231,
274
Kippistia 126, 130, 133, 134, 144,
147, 210, 234, 264, 274
Knozia 463
 sumatrensis 463
Krylovia 201, 273

Labiatae 280, 382
Laburnum 441, 442, 444, 445
 anagyroides 444, 445
Lachnophyllum 205, 223, 273
Lactoridaceae 3
Lactoridales 3
Laennecia 101, 103, 115, 123-127,
209, 210, 234, 235, 250, 266,
272
 filaginoides 235
 gnaphalioides 235
 mapimiana 123
 sophiiifolia 235
 schiedeana 235
Laestadia 90, 207, 225, 228, 273
Lagenifera 136, 137, 141, 143, 148,
151-153, 155, 156, 158, 197,
208, 230, 231, 251, 261, 274
 huegelii 155
 sect. *Lagenifera* 261
 sect. *Pseudomyriactis* 230
 stipitata 155
Lageniferinae 193, 198, 207, 215,

- Lageniferinae (cont.) 225, 230-232,
 245, 246, 248, 251, 256, 273,
 274
Lagenithrix 143, **148-152**, 155, 156,
 208, 230, 231, 274
 setosa 143, 148-150, 151
 stellata 143, 148, 149, **151**, 152
Lagenopappus 143, 149, 152, **153-**
 156, 208, 230, 231, 274, 426
 gunnii 143, 149, 153, **154**, 155,
 426
 pappocromus 143, 153, **154**,
 155
 tasmanicus 143, **154**, 155, 426
Lagenophora 158, 260
 Lamiaceae 277, 345, 382, 391, 463,
 476
Lamiastrum 454
 Lamiidae 3, 5
Lantana 464
 camara 464
Laphamia 324, 327, 328
 ambrosiaefolia 324
 ambrosifolia 327
Lasianthaea 40
Lasiocephalus 402, 404
 stylotrichus **404**
 Lauraceae 454
 Leguminosae 302, 384, 388, 420, 425
 Lentibulariaceae 69
Lepidaploa 27-29
 balansae **27**
 marguana **28**
 paraensis **28**
 pellita **28**
 spixiana **28**
Lepidophyllum 207, 273
Leptoglottis 414, 417, 421, 422, 424,
 425
 hystricina 414
 latidens 422
 nelsonii 414, 421, 425
 nuttallii 417
Leptoglottis (cont.)
 occidentalis 422
 paucijuga 424
 potosina 422, 424
 regiomontanus 424
Leptostelma 122, 124, 159, 167-170,
 174-176, 182-184, 198, 204,
 220, 248, 267, 273
 tweediei 183
Lessingia 209, 232-234, 264, 265, 272
Lessingianthus 27-29
 regis 28
 scabrifolius **29**
Leucas 463
 aspera 463
Leucelene 271
Leucopholis 185, 188
Leucophyllum 386
 hintoniorum 386
Leucopsis 111, 115, 117
 podocomoides 111
 scaposa 117
Leucosyris 272
Leyssera 92
 ciliata 92
 Liabeae 19, 23
 Liliidae 2
 Liliopsida 1, 2
 Limnanthaceae 4
 Limnanthales 4
Lindernia 463
 crustacea 463
Linosyris 273
Lippia 464
 nodiflora 464
Litsea 367
Llerasia 207, 273
Lucilia 187, 229
Lupinus 290-300, 321, 510
 bartlettianus 290, 292, 293, 296
 caballoanus 290, 291, **294-296**,
 300
 cacuminis 291, 295, 296

- Lupinus* (cont.)
elegans 300
harleyanus 293
havardii 290, 291, 295, 298
hintonii 299
hintoniorum 290, 291, 295, 296, 299, 300
leonensis 290, 301, 302
montanus 290, 295, 302
muelleri 290, 292, 295, 297, 299, 300
oligostolus 293
omalophyllus 293
 "partisans" [should have been *potosinus*] 290, 292, 294, 297, 299, 300, 510
 var. *bartlettii* 292
platamodes 290, 292, 294, 297, 300
polyphyllus 321
potosinus 290, 292, 294, 295, 297, 299, 300, 510
stipulatus 290, 291, 297, 300, 301
 var. *exilis* 301
stolidotus 293
subcarnosus 301, 302
tamaulipensis 293
tezensis 290, 291, 295, 298, 301, 302
- Lycium* 362
Lythrum 369
- Machaeranthera* 194, 198, 208, 209, 232, 233, 240, 263, 265, 272
 sect. *Psilactis* 265
- Machaerantherinae* 193, 198, 199, 208, 232, 233, 238, 245, 249, 250, 271-273
- Macronema* 272
- Madagaster* 90, 95, 207, 247, 267, 274
- Magnoliales 3
 Magnoliidae 2, 6, 7
- Magnoliophyta 1-3, 7
 Magnoliopsida 1-3
Mairia 85-88, 90-94, 198, 199, 207, 222, 228, 267, 274
burchellii 91
coriacea 91
corymbosa 85, 88, 93
crenata 86, 90, 91
decumbens 91
ecklonis 91
felicioides 94
foliosa 92
hirsuta 91
 sect. *Homochroma* 91
lasiocarpa 92
microcephala 92
montana 92
perezioides 86, 92
pinifolia 92
 sect. *Pteropappus* 91
pustulata 94
tazifolia 92
 sect. *Zyrphelis* 91
- Malvaceae 307, 308, 322, 463
Martynia 463
annua 463
Matricaria 474
matricarioides 474
Melanodendron 202, 213, 214, 247, 274
Melhania 464
cannabina 464
Merope 403
aretioides 403
Merremia 462
emarginata 462
tridentata 462
Mesoligus 272
Microcalia 274
Micrococca 463
mercurialis 463
Microglossa 202, 213, 274
Microgyne 101, 102, 104

- Microgyne* (cont.)
trifurcata 104
- Microgynella* 101, 103, 104, 114, 115, 122, 124, 126, 175, 210, 234, 235, 267, 273
trifurcata 104
- Microtrichia* 274
- Mimosa* 412-425, 477
 sect. *Batocaulon* 413
biuncifera 477
hystricina 412-414, 415, 417
latidens 412, 414, 416-418, 421-425
microphylla 412, 413, 418
 ser. *Mimosa* 413
 sect. *Mimosa* 413
nutallii 412-415, 417, 424
occidentalis 412, 413, 416, 417, 421-423, 425
paucijuga 421, 422-424
platycarpa 417
potosina 421-424, 425
 ser. *Quadrivalves* 413, 425
quadrivalvis 412, 414, 417, 418, 418, 420-422, 424, 425
 var. *angustata* 418
 var. *latidens* 422, 424
 var. *nelsonii* 421, 425
 var. *paucijuga* 424
 var. *platycarpa* 418
roemeriana 412, 413, 417-419
subinermis 421-424, 425
- Mimosaceae 412, 421, 425
- Mimosoideae 420, 425
- Minuria* 125, 126, 130, 133, 134, 144, 146, 147, 157, 197, 200, 210, 216, 218, 227, 234, 237, 261, 264, 274
denticulata 157
integerrima 261
- Miyamayomena* 201, 273
- Mniodes* 187, 403, 405
andina 403
- Mniodes* (cont.)
aretioides 403
pulvinulata 403
- Mnioides* 229
- Mollugo* 462
nudicaulis 462
pentaphylla 462
- Monogyria* 170
- Monoptilon* 88, 199, 206, 223-225, 250, 251, 272
- Moraceae 477
- Moringaceae 4
- Moringales 4
- Moritzia* 112
- Morongia* 414, 417, 422
latidens 414, 422
occidentalis 417, 422
- Morus* 477
alba 477
- Mukia* 462
scabrella 462
- Munnozia* 19-23
chachapoyensis 19, 20, 21
hastifolia 22
luyensis 19, 21, 22
perfoliata 21
senecionidis 22
smithii 19, 22, 23
subviridis 22
- Mutisiaeae 27, 262
- Myoporaceae 246
- Myriactis* 141, 148, 151, 152, 155, 156, 208, 230, 231, 274
- Nannoglottis* 211, 237-240, 265, 273
- Nardophyllum* 199, 207, 228, 251, 267, 273
scoparium 267
- Navarreta* 305
- Navarretia* 56
incisa 56
- Neja* 101-105, 122, 124, 168-173, 176, 182-184, 198, 204, 220, 273

- Neja* (cont.)
ciliaris 171
dianthifolia 168, **171**, 172
falcata 172
filiformis 103, 169-172
gracilis 170, 171
linearifolia 170, 172
macrocephala 104, 172
marginata 168-171
montevidensis 171
nidorelloides 169, 171
sect. *Phylloneja* 104, 172
pinifolia 168-171, 173
sect. *Podoneja* 170
pulvinata 168, 169 **172**
sprengelii 172
subvillosa 172
tenuifolia 171
- Neomolina* 214, 263, 273
Neopazia 150
Neurotheca 8
Nidorella 197, 198, 206, 222, 227,
228, 247, 271, 274
undulata 227
- Niebla* 458-460
cedrosensis 458-460
homalea 458
josecuervoi 460
- Nolletia* 87, 197, 205, 274
Noticastrum 115, 124, 198, 204, 220,
249, 250, 271, 273
Novenia 207, 228, 229, 262, 273
acaulis 229
- Nyctanthaceae 344
- Oblivia* 24-26
ceronii **24**
mikanioides 25
simplex 25
Ocimum 463
americanum 463
basilicum 463
- Oclemena* 212, 272
- Olearia* 130, 163, 165, 166, 195, 197,
207, 228, 230, 242, 243, 247,
251, 261, 274
chathamica 165
colensoi 165
lyalii 165
oporina 165
rapa 163
- Oldenlandia* 463
corymbosa 463
herbacea 463
umbellata 463
- Oligandra* 191
Oliganthes 18
jelskii 18
- Oligoneuron* 211, 237, 266, 272
Olivaea 209, 233, 272
Onopordum 474
acanthium 474
- Oonopsis* 209, 232, 272
Operanthus 163
Opuntia 37
- Oreochrysum* 211, 237, 239, 240, 272
Oreostemma 212, 272
- Oritrophium* 90, 121, 123, 167, 207,
227, 228, 266, 273
orizabense 123, 167, 266
- Osbertia* 116, 204, 272
Osteospermum 93
falcatum 93
- Ozothamnus* 188
- Pachystegia* 165, 207, 274
- Pacifigeron** 160, 161, 163-166, 197,
207, 228, 267, 274
rapensis 160, 161, **163**, 164
- Paleaepappus* 228, 273
Panicum 463
repens 463
- Papaveraceae 463
Pappochroma 153, 426
gunnii **426**
tasmanica **426**

- Pappochroma* (cont.)
uniflora 153, 426
Parachionolaena 185, 191
Parastrephia 207, 267, 273
Parnassiales 4
Parthenium 462
hysterophorus 462
Pavonia 463
zeylanica 463
Pedaliaceae 463
Pedaliium 463
murex 463
Pediculareae 409
Pentacalia 402, 404
andicola 404
aquifolia 404
curvidens 404
guadalupe 404
ilicifolia 404
subg. *Microchaetae* 402, 404
miniaurita 404
pavonii 404
pleniaurita 404
Pentachaeta 198, 200, 206, 224, 225,
250, 270, 272
Pentadiplandraceae 4
Pergularia 462
aemia 462
Peripleura 124, 125, 131-133, 144,
146, 147, 159, 210, 237, 267,
274
arida 131
bicolor 131
diffusa 131
hispidula 131
var. *hispidula* 131
var. *setosa* 131
obovata 131
scabra 131
sericea 132
spechtii 132
Peripleura spechtii (cont.)
var. *kimberleyensis* 132
var. *spechtii* 132
virgata 132
Peritylanae 328
Perityle 324-328
ambrosiifolia 324, 325-327
coronopifolia 327
dissecta 327
gilensis 324, 327
sect. *Laphamia* 328
lemmonii 324, 327
sect. *Perityle* 328
sazicola 324, 327
Perotis 463
indica 463
Petalacte 187
Petradoria 237, 238, 272
Phacellothrix 242
cladochaeta 242
Phyllanthus 463
amarus 463
debilis 463
maderaspatensis 463
virgatus 463
Phyllostelidium 402, 403
aretioides 402
aretioideum 402, 403
pavonianum 402, 403
Physalis 441, 442, 452-454, 457, 464
izocarpa 441, 452
var. *immaculata* 441, 452
var. *parviflora* 441, 452
minima 464
philadelphica 441, 452, 454
var. *immaculata* 452
var. *parviflora* 452
var. *philadelphica* 452
Pinaceae 73
Pingraea 214, 263, 273
Pinguicula 69-72

- Pinguicula* (cont.)
esseriana 69, 71, 72
hintoniorum 69, 71
jorgehintonii 69-72
- Pinus* 73-79, 475, 478
 subg. *Balfouria* 73, 77, 78
Diplozylon 73
 subg. *Ducampopinus* 78
edulis 475, 478
 subg. *Gerardia* 73, 77, 78
Haplozylon 73
nelsonii 77
 subg. *Pinea* 73, 74, 77, 78
 subg. *Pinus* 73, 77, 78
ponderosa 478
rzedowskii 73
 subg. *Sabinia* 73, 74, 77, 78
strobis 74, 76
 subg. *Strobis* 73, 77, 78
sylvestris 74, 75
 subg. **Tamaulipasa** 73, 77, 78
- Piora* 151, 156, 208, 231, 274
- Piperaceae 3, 7
- Piperales 3, 7
- Piperanae** 1, 3
- Piperidae** 1, 2, 3
- Piperinae 3
- Piperopsida 1-3
- Pittosporales 5
- Pityopsis* 204, 269, 272
- Plagiocheilus* 200, 206, 225, 269, 273
- Plantago* 150
- Platanaceae 275
- Platanus* 250
- Pleurophyllum* 165, 166, 207, 261, 274
- Plucheae 259
- Plumbaginaceae 30, 31
- Plumbago* 30-36
scandens 30-36
- Poa* 159
- Poaceae 410, 411, 463
- Podocoma* 101, 103-113, 115, 121, 122, 124, 126, 128, 170, 209, 210, 234, 235, 268, 273
asperrima 110, 111
bartsiiifolia 106, 112
bellidifolia 107-109, 111
blanchetiana 109, 111
erigerifolia 111
foliosa 111
hieracifolia 108, 109, 111
hirsuta 109-111
macrocephala 104, 106, 112
notobellidiastrum 106, 108, 110, 112
primulifolia 111
regnellii 110, 112
reineckii 113
rivularis 106, 108, 110, 112
spegazzinii 111
- Podocominae** 193, 197, 198, 209, 218, 227, 230, 234-237, 242, 246, 248, 250, 255, 272, 274
- Podopappus* 104, 106, 109, 111
hirsutus 109, 111
pubescens 111
tomentosus 104
- Poecilolepis* 197, 205, 223, 274
- Polemoniaceae 52, 68, 69, 303, 305, 477
- Polemonium* 56
incisum 56
- Polyactidium* 171
sprengelii 171, 172
- Polyarrhena* 85, 87-89, 93, 94, 197, 205, 262, 274
refleza 89
 subsp. *refleza* 89
- Populus* 250, 477
fremontii 477

- Portulaca* 463
 oleracea 463
 quadrifida 463
 Portulacaceae 463
Prionopsis 232, 268, 272
Prosopis 325
Prunus 441, 442, 451, 454
 cuneata 451
 pumila 441, 451
 var. *cuneata* 441, 451
 var. *susquehanae* 441, 451
 susquehanae 451
Psednotrichia 244
 australis 244
 tenella 244
Pseudobaccharis 273
Pseudolarix 78
Pseudoligandra 185, 191
Pseudolinosyris 273
Psiadia 202, 213, 247, 274
Psiadiella 202, 274
Psilactis 212, 240, 250, 272
 Psilotaceae 275
Psychrogeton 198, 201, 273
 Pteridophyta 142
Pteronia 90, 197, 207, 222, 274
Pyrrocoma 209, 232, 238, 272
Pytinicarpa 136-142, 159, 208, 230,
 232, 267, 274
 neocaledonica 136, 138, 140
 sarasinii 136, 138, 139, 141,
 142
Quercus 310, 367, 370, 475, 476, 478
 arizonica 475
 emoryi 475
 gambelii 478
 havardii 476
Racinaea 284-289
 adpressa 286
 Racinaea (cont.)
 blassii 286
 euryelytra 284-286
 flexuosa 289
 kalliantha 284, 286, 287
 lymansmithiana 284, 286, 288
 parviflora 284, 286
Ramalina 460
 Ramalinaceae 458
 Ranunculaceae 442, 445, 456
 Ranunculales 4
 Ranunculidae 2
 Ranunculopsida 1, 2
Ranunculus 441, 442, 446, 447, 476
 glaberrimus 441, 446, 447
 var. *reconditus* 441, 446, 447
 reconditus 446, 447
 testiculatus 476
 triternatus 446, 447
Raouliopsis 187
Remya 197, 207, 228, 270, 274
 Resedaceae 4
Rhabdanthus 101, 103, 236
 Rhamnaceae 442, 447, 454, 457
 Rhamnales 454
Rhamnus 250, 441, 442, 447-451, 453,
 454, 456, 457
 betulifolia 448
 var. *obovata* 448
 blumeri 448
 californica 448, 449
 ssp. *crassifolia* 448
 var. *crassifolia* 448
 ssp. *cuspidata* 449
 ssp. *occidentalis* 449
 var. *occidentalis* 449
 ssp. *tomentella* 449
 ssp. *ursina* 449
 var. *ursina* 449
 cathartica 450

- Rhamnus cathartica* (cont.)
 var. *davurica* 450
 citrifolia 441, 450, 456
 var. *nipponica* 450
 cuspidata 449
 davurica 441, 450
 ssp. **nipponica** 441, **450**
 var. *nipponica* 450
 lanceolata 441, 450, 451
 ssp. **glabrata** 441, **451**
 var. *glabrata* 450, 451
 nevadensis 449
 obtusissima 449
 occidentalis 449
 rubra 449
 ssp. *modocensis* 449
 ssp. *nevadensis* 449
 ssp. *obtusissima* 449
 ssp. *yosemitana* 449
 sphaerosperma 450
 var. *longipes* 450
 tomentella 448, 449
 ssp. *crassifolia* 448
 ssp. *cuspidata* 449
 ssp. *ursinus* 449
 ursina 449
Rhamphogyne 208, 230, 231, 274
Rhus 250
Rhynchosia 463
 minima 463
 suaveolens 463
Rhynchospermum 208, 215, 231, 274
Rigiopappus 206, 224, 225, 251, 268,
 269, 272
Rochonia 90, 198, 207, 274
Rosaceae 442, 451, 457
Rosales 4
Rosanae 4
Rosidae 3-5
Rosopsida 1, 3
Rubiaceae 463, 477
Rubus 310
Ruellia 461
 tuberosa 461
Rungia 461
 pectinata 461
 repens 461
Rutanae 4
Rutidosia 242
 brownii 242
Salvadoraceae 4
Salvadorales 4
Salvia 476
 pratensis 476
Sapindaceae 463
Sapindanae 4
Sarcanthemum 202, 274
Saururaceae 3, 457
Saxifragaceae 4
Saxifragales 4
Saxifraganae 1, 4
Schizoptera 39, 50
 peduncularis 50
 trichotoma 50
Schrankia 412-414, 418, 421, 422,
 424, 425
 hystericina 412, 414
 latidens 412, 414, 422, 425
 microphylla 412, 418
 nuttallii 414
 occidentalis 412, 422
 platycarpa 418
 potosina 414
 quadrivalvis 422
 var. *occidentalis* 422
 roemeriana 412
 subinermis 424, 425
 uncinata 412
Scoparia 463
 dulcis 463
Scorzonera 474, 478

Scorzonera (cont.)*laciniata* 474, 478*Scrobicaria* 402, 404*aquifolia* 404*ilicifolia* 404*Scrophulariaceae* 329, 332, 406, 409,
463, 465, 467*Scutellaria* 345-382*alata* 350, 352, 354*alpina* 381*angustifolia* 382*anomala* 345, 350, 352, 354*aramberrana* 345, 351, **352**, 354*bartlettii* 345, 347, 349, **355**,
356, 361*brevifolia* 380*carmenensis* 345, 346, 351, 356,
361*chiangii* 345, 350, 357, **361***drummondii* 345-347, 349, 353,
355, 362-365, 370, 379-381var. *drummondii* 345, 353, 363-
365, 370var. *edwardsiana* 345, 353,
362, **364**, 365, 379, 381var. *runyonii* 345, 353, 363,
365*durangensis* 345, 350, 356, **365**-
367*frazinea* 366, 379*fruticetorum* 345, 347, 351, 356,
366sect. *Galericularia* 345, 347, 366,
379*helleri* 363*hispidula* 345, 351, 357, 362, 366,
367, 378*horridula* 345, 367*incertis* 370*integrifolia* 380*Scutellaria integrifolia* (cont.)var. *brevifolia* 380*laevis* 345, 346, 349, 358, 367,
368sect. *Lateriflorae* 347, 366, 379*lythroides* 369*melanquitensis* 346*microphylla* 361, 368, 369, 380,
381sect. *Mixtae* 366*monterreyana* 345, 351, 355,
356, 361, **368**, 380*mulleri* 345, 351, 357, 368, **369***muriculata* 345, 349, 358, 365,
369, 370*muzquiziana* 345, 347, 351, 357,
361, 369, **370**, 371*molanquitensis* 381*potosina* 345, 346, 349, 351, 352,
355, 359, 361, 362, 368, 370-
377, 381var. *davisiana* 345, 359, 372,
373, 374, 376, 377var. *grahamiana* 345, 359, 368,
372, **374**var. *novoleonensis* 345, 359,
372-**374**, 375, 376subsp. *parviflora* 374, 376subsp. *platyphylla* 370, 371, 374,
376, 377var. *platyphylla* 345, 359, 370,
376, **377**subsp. *potosina* 371-377var. *potosina* 345, 352, 359, 361,
372, 373, 375-377var. *tessellata* 345, 352, 359,
371-**375**, 376, 377var. *zacatecensis* 345, 359,
372, **376***resinosa* 345, 348, 349, 358, 364,

- Scutellaria resinosa* (cont.) 365, 377, 380, 381
 var. *brevifolia* 380
 var. *resinosa* 381
 sect. *Resinosa* 345-348, 366, 379-381
 sect. *Spinosa* 346, 381
stewartii 345, 347, 350, 352, 354, 367, **378**, 379
tessellata 375-377
texana 345, 347, 349, 350, 356, 361, 368, **379**-381
thieretii 346, 363
wrightii 345, 347, 349, 350, 360, 362, 365, 365, 376, 377, 380, 381
- Scutellarioideae 346
Sebastiania 463
chamaelea 463
- Senecio* 404
curvidens 404
miniauritus 404
pleniauritus 404
scrobicaria 404
stylotrichus 404
- Senecioideae 402
 Senecioneae 238, 239, 248, 264, 404
Sericocarpus 212, 240, 250, 272
Sheareria 208, 231, 273
- Sida* 463
acuta 463
cordifolia 463
rhombifolia 463
veronicifolia 463
- Sidalcea* 307-316, 321-323
nelsoniana 307-316, 321-323
- Sideranthus* 272
- Sisyrinchium* 468-472
arizonicum 471
cernuum 469
convolutum 471
- Sisyrinchium* (cont.)
conzattii 471
 subg. *Echthronema* 468, 471
hintoniorum **468**, 470, 471
longipes 469
microbracteatum 468, **470**, 471
palmeri 471
schaffneri 468-470, 472
 subg. *Sisyrinchium* 472
tenuifolium 468, 470, 471
tolucense 471
- Solanaceae 362, 442, 452, 454, 464
Solanum 464
nigrum 464
surattense 464
trilobatum 464
- Solenogyne* 141, 151, 152, 156, 158, 208, 230, 231, 259, 261, 274
- Solidagininae 193, 194, 198, 201, 210, 211, 237-241, 245, 246, 249, 250, 271, 272
- Solidago* 193, 195, 197, 198, 200, 210, 211, 237, 239, 240, 250, 266, 272
- Sommerfeltia* 101-105, 114, 115, 122, 124, 126, 127, 175, 197, 210, 234-236, 267, 273
cabreræ 103, 235, 236
spinulosa 104, 236
- Sonchus* 462
oleraceus 462
- Sophora* 385-388
arizonica 385
Calia grouping 385
formosa 385
gypsophila 385-387
hintoniorum 386
juanhintoniana **385**-387
purpusii 385-387
- Sophoreae 387

- Spermatophyta 451
- Sphaeranthus* 462
indicus 462
- Sporobolus* 329, 330, 474
airoides 329, 330, 474
- Stachys* 391-401
albotomentosa 391-394, 400
 var. *albotomentosa* 391, 393, 394, 400
 var. *potosina* 391, **393**, 400
cardinalis 394
coccinea 391, 392, 394-398, 401
jaimehintonii 391, 392, **395**, 400
limitanea 394
lindenii 391, 392, 396, 399, 401
oaxacana 394, 395
pacifica 391, 392, 395, **396**, 397, 400
torresii 391, 392, **397**, 400, 401
- Stachelina* 404
ilicifolia 404
- Stemodia* 463
viscosa 463
- Stenachaenium* 113
campestre 113
- Stenandrium* 433-439
barbatum 433-437, 439
chameranthemoideum 433-437, 439
droseroides 434
dulce 433-437, 439
manchonense 433-437, 439
nanum 433-435, 437-439
pedunculatum 433-435, 437-439
pilosulum 433, 434, 437-439
pohlii 434
subcordatum 434, 435
trinerve 434
verticillatum 434, 435
- Stenotopsis* 272
- Stenotus* 211, 238, 272
- Stephanodoria* 209, 272
- Sterculiaceae 464
- Stereosanthus* 238
- Steviopsis* 389, 390
fendleri 389, **390**
nesomii 389, 390
- Stilbaceae 344
- Stipa* 475
comata 475
- Streptoglossa* 157
decurrens 157
liatroides 157
- Striga* 463
densiflora 463
- Stylidiaceae 246
- Styrax* 250
- Symphoremaceae 344
- Symphyotrichinae** 193, 194, 197, 198, **211**, 212, 237, 239-241, 245, 249-251, 271, 272
- Symphyotrichum* 200, 212, 221, 239, 240, 250, 251, 272
 subg. *Brachyactis* 221, 251
- Synapsis* 443
recurvata 443
- Synedrella* 39-46, 51
nodiflora 39, 42-46
peduncularis 39, 51
- Synedrellopsis* 39
- Tageteae 268
- Tamarix* 477
ramosissima 477
- Tephrosia* 463
hirta 463
procumbens 463
purpurea 463
- Tetramelaceae 4
- Tetramolium* 125, 126, 128-134, 144, 197, 209, 210, 234, 265, 274

- Tetramolopium* (cont).
 sect. *Alpinum* 129, 130
bicolor 130
humile 129, 130
 sect. *Sandwicense* 129, 130
sylvae 129
 sect. *Tetramolopium* 129, 130
- Themeda* 463
triandra 463
- Thermopsis* 383, 384
gracilis 383
 var. *ovata* **383**
macrophylla 384
 var. *hitchcockii* 384
montana 383, 384
 var. *hitchcockii* 383, **384**
 subsp. *ovata* 383
 var. *ovata* 383
ovata 383
rhubifolia 383
 var. *ovata* 383
- Thespis* 208, 231, 274
- Thurovia* 211, 272
- Tiliaceae 464
- Tillandsia* 286
adpressa 286
- Tillandsioideae 284
- Tolbonia* 217, 218, 259, 273
anamitica 217, 218
- Tomentaurum* 204, 272
- Tonestus* 212, 237, 240, 266, 268,
 272
kingii 240
- Toricelliales 5
- Tovariaceae 4
- Townsendia* 193, 198, 203, 216, 219,
 223, 251, 259, 265, 271, 272
ezimia 219
formosa 219, 251
microcephala 261
- Tracyina* 206, 224, 268, 272
- Trianthes* 462
portulacastrum 462
- Tribulus* 464
terrestris 464
- Trichilia* 367
- Trichodesma* 462
indicum 462
- Tridax* 462
procumbens 462
- Trimeniaceae 3
- Trimorpha* 221, 222, 265, 272
acris 222
- Tripolium* 201, 273
- Triumfetta* 464
pentandra 464
rhomboidea 464
- Triurididae 2
- Tubuliflorae 105, 123, 183, 263
- Turczaninowia* 273
- Tylophora* 462
indica 462
- Ucacou* 46
nodiflorum 46
- Ulmus* 477
pumila 477
- Umbraculariae 4
- Unamia* 272
- Uncinia* 159
- Urena* 463
lobata 463
- Urochloa* 463
panicoides 463
- Vancklevea* 211, 238, 259, 272
- Verbenaceae 344, 464
- Verbesina* 39, 46, 281-283
microptera 281, 282
nodiflora 39, 46
 sect. *Ochractinia* 281-283

Verbesina (cont.)*rumicifolia* 282**tamaulipana** 281, 282*virginica* 281-283*Vernonanthura* 27, 29*patens* 29**santacruzensis** 29*Vernonia* 27-29, 462*aristeguietae* 27*balansae* 27*cinerea* 462*marguana* 28*paraensis* 28*pellita* 28*santacruzensis* 29*scabrifoliata* 29*sparrei* 27*spiziana* 28*subcordata* 28*trichotoma* 27*velascens* 29*Vernonieae* 18, 26, 27, 29*Vernoniopsis* 202, 243, 274

Viburnales 5

Vicia 312*Vierhapperia* 238*Viola* 478*nuttallii* 478

Violaceae 478

Violales 3, 4

Violanae 3, 4

Virgulus 272*Vittadinia* 88, 101, 102, 104, 105,

113, 122, 124-134, 143, 144,

146, 147, 159, 195, 199, 210,

225, 260, 267, 274

arida 131*australis* 157*bicolor* 131*cervicularis* 127*Vittadinia* (cont.)*diffusa* 131*hispidula* 127, 131var. *setosa* 131*macrorrhiza* 125*megacephala* 127, 128*obovata* 131subg. *Peripleura* 125-129, 131*pterochaeta* 127, 128*pustulata* 127, 128*scabra* 127, 131*sericea* 132*spechtii* 132var. *kimberleyensis* 132*trifurcata* 104*virgata* 132subg. *Vittadinia* 125-129, 131*Waltheria* 464*americana* 464*Wamalchitamia* 26*Wardaster* 273*Weberaster* 272*Wedelia* 26, 40, 191*Westoniella* 207, 228, 261, 272*Wyomingia* 272*Xanthisma* 197, 209, 272*Xanthium* 462*strumarium* 462*Xanthocephalum* 209, 232, 272*Xiphochaeta* 28*Xylorhiza* 198, 209, 232, 238, 272*Xylothamia* 198, 211, 268, 272*Yucca* 475*elata* 475*Zalazaria* 346

Zalazariaceae 346

Zehneria 444*cerasiformis* 444

- Zexmenia* 26
 mikanioides 26
- Zingiberidae 2
- Zornia* 463
 diphylla 463
- Zygophyllaceae 464
- Zyrphelis* 85-94, 197-199, 205, 211,
 222, 224, 225, 274
 amoena 91, 92
 angolensis 85, 89, **91**
 burchellii 91
 corymbosa 93
 crenata 91
 decumbens 85, **91**
 ecklonis 89, 91
 foliosa 92
 hirsuta 91
 lasiocarpa 89, 92
 microcephala 92
 montana 85, **92**
 perezoides 85, 87, 91, **92**
 tazifolia 89, 91, 92
- Zyzyzia* 26

INDEX TO REVIEWERS, VOLUME 76

The editor express his most sincere appreciation to the following individuals. These are persons who have reviewed papers that were submitted for publication in volume 76 of *Phytologia*. Without the willingness and diligence of these reviewers, the task of the editor would be much more difficult, and the quality of the papers published would be lessened. To each of you, I offer my most sincere thanks.

Michael J. Warnock, Editor.

- | | |
|-------------------|-------------------|
| Allred, K. | McLaughlin, S. |
| Barkworth, M.E. | Nesom, G.L. |
| Bowler, P.A. | Prather, A. |
| Brown, L.E. | Pruski, J. |
| Cayouette, J. | Ramamoorthy, T.P. |
| Chilcote, W.W. | Reveal, J.L. |
| Cody, W.J. | Robinson, H. |
| Day, A. | Rodríguez J., C. |
| Delprete, P. | Rundell, J.R. |
| Dillon, M. | Spellenberg, R.M. |
| Dutton, B.E. | Strother, J. |
| Egger, M. | Turner, B.L. |
| Fernández, N., R. | VanDevender, R. |
| Fishbein, M. | Wasshausen, D.C. |
| Grant, V. | Watson, T. |
| Jansen, B. | Wieboldt, T.F. |
| Johnston, L.R.D. | Wiersema, J.H. |
| Karis, P.O. | Zanoni, T. |
| Lellinger, D.B. | |
| Levin, G. | |
| Linsley, E.G. | |
| Luther, H.E. | |
| Mason, C.T. | |
| Mayfield, M. | |

CORRECTIONS AND ADDITIONS

Volume 74, issue 3, page 178, line 4 of abstract, "Fab-anae" should read "Fabanae".

Volume 75, issue 2, inside front cover, contents, "SUTHERLAND, D.M." omitted as an author on the paper entitled "Chromosome numbers for *Dalea* species (Fabaceae) from southwestern New Mexico and southeastern Arizona.

Volume 76, issue 4, on pages 290-302, due to an editorial error made during spell checking the manuscript, the specific epithet *potosinus* was replaced with "partisans" throughout the manuscript.

AVAILABILITY OF BACK ISSUES OF PHYTOLOGIA

The following back issues of PHYTOLOGIA are available:

37(3)	September 1977	\$3.00
41(2,3,6,7)	December 1978-March 1979	\$3.00 each
42(1)	April 1979	\$3.00
43(1-3)	May-August 1979	\$3.00 each
44(2,4,7)	August-December 1979	\$3.00 each
45(1,3-6)	February-June 1980	\$3.00 each
46(1,5-7)	June-October 1980	\$3.00 each
47(1,2,4,6)	November 1980-March 1981	\$3.00 each
48(1,3-6)	April-August 1981	\$3.00 each
49(1-5)	September-November 1981	\$3.00 each
50(1-7)	December 1981-May 1982	\$3.00/issue or \$17.00 for complete volume
51(1-7)	May-September 1982	\$3.00/issue or \$17.00 for complete volume
52(1-7)	September 1982-March 1983	\$3.00/issue or \$17.00 for complete volume
53(1-7)	March-August 1983	\$3.00/issue or \$17.00 for complete volume
54(1-7)	September 1983-January 1984	\$3.00/issue or \$17.00 for complete volume
55(1-7)	February-June 1984	\$3.00/issue or \$17.00 for complete volume
56(1-7)	July 1984-January 1985	\$3.00/issue or \$17.00 for complete volume
57(1-7)	February-August 1985	\$3.00/issue or \$17.00 for complete volume
58(1-7)	August-December 1985	\$3.00/issue or \$17.00 for complete volume
59(1-7)	December 1985-April 1986	\$3.00/issue or \$17.00 for complete volume
60(1-4,6,7)	May-September 1986	\$3.00/issue or \$17.00 for those available
61(1-7)	September 1986-January 1987	\$3.00/issue or \$17.00 for complete volume
62(1-6)	February-May 1987	\$3.00/issue or \$17.00 for complete volume
63(1-6)	May-November 1987	\$3.00/issue or \$17.00 for complete volume
64(1-6)	December 1987-June 1988	\$3.00/issue or \$17.00 for complete volume
65(1-6)	June 1988-February 1989	\$3.00/issue or \$17.00 for complete volume
66(1-6)	April-July 1989	\$3.00/issue or \$17.00 for complete volume
67(1-6)	August-December 1989	\$3.50/issue or \$21.00 for complete volume
68(1-6)	January-June 1990	\$3.50/issue or \$21.00 for complete volume
69(1-6)	August-December 1990	\$3.50/issue or \$21.00 for complete volume
70(1-6)	January-June 1991	\$3.50/issue or \$21.00 for complete volume
71(1-6)	August-December 1991	\$3.50/issue or \$21.00 for complete volume
72(1-6)	January-June 1992	\$3.50/issue or \$21.00 for complete volume
73(1-6)	August-December 1992	\$3.50/issue or \$21.00 for complete volume

Postage: \$.50 per issue domestic, \$2.00 per volume domestic; \$1.00 per issue foreign, \$4.00 per volume foreign.

Payable in U.S. dollars free of any bank charges to: PHYTOLOGIA, 185 Westridge Drive, Huntsville, TX 77340-8916 U.S.A.

Telephone: 409 295-5410; FAX: 409 291-0009; Federal EIN: 76-0274455.

Inquire on discounts for large orders (more than five volumes).

All back issues are subject to prior sale.

New York Botanical Garden Library



3 5185 00288 4755

Information for Authors

Articles from botanical systematics and ecology, including biographical sketches, critical reviews, and summaries of literature will be considered for publication in PHYTOLOGIA. Manuscripts may be submitted either on computer diskette, or as typescript. Diskettes will be returned to authors after action has been taken on the manuscript. Diskettes may be 5.25 inches or 3.5 inches and may be written in any IBM or MacIntosh compatible format. Typescript manuscripts should be single spaced and will be read into the computer using a page scanner. The scanner will read standard typewriter fonts but will not read dot matrix print. Manuscripts submitted in dot matrix print cannot be accepted. Use underscore (not italics) for scientific names. Corrections made on typescript manuscripts must be complete and neat as the scanner will not read them otherwise. Language of manuscripts may be either English or Spanish. Figures will be reduced to fit within limits of text pages and therefore, should be submitted with an internal scale and have dimensions proportional to those for text pages. Legends for figures should be included in figures whenever possible. Each manuscript should have an abstract and key word list. Specimen citations should be consistent throughout the manuscript. Serial titles should be cited with abbreviations used in *Botanico Periodicum Huntianum*. References cited only as part of nomenclatural summaries should not appear in Literature Cited. Nomenclatural work should include one paragraph per basionym and must provide proper (as defined by the current *International Code of Botanical Nomenclature*) citation of sources of epithets and combinations.

Authors should arrange for two workers in the appropriate field to review the manuscript before submission. Copies of reviews should be forwarded to the editor with the manuscript. Manuscripts will not be published without review.

Cost of publication is currently \$13.00 US per page for publication without reprints. Publication with 100 reprints is provided for \$18.00 US per page, 200 reprints for \$21.50 US per page. Page charges are due with manuscript and no paper will be published before payment is received in full. Reprints must be ordered and paid for in advance. Page charges will be determined on the basis of a typescript page (single spaced, 10 points, blank line between paragraphs) with all type inside a rectangle 143 mm (horizontal) by 219 mm (vertical), not including running head and page number. Title page should include title, author(s) name(s), and address(es). Two blank lines should appear above and below section headings (Abstract, Discussion, Literature Cited, etc.) in the manuscript. No extra charge is made for line drawings provided they conform to limitations of size and proportion for normal text. Halftones require an extra charge of \$10.00 US per page.